## The Butterflies and Moths of Leicestershire and Rutland

## Provisional List of Macromoths Recorded from VC55

## 2016 Update - version 2

This is a provisional list that excludes a number of species only doubtfully recorded from the county (some of which were recorded in Victoria County History lists and also excludes a number species where records are still under investigation.

With work continuing towards publication of "The Butterflies and Moths of Leicestershire & Rutland", the aim of this document is to provide local recorders with an initial indication of the current state of knowledge as far as the status and distribution of macromoths in the County is concerned. Hopefully, this will highlight some of the research still required, as well as stimulating further fieldwork which could help to answer many of the outstanding uncertainties surrounding the County's macro-moths.

However, at this stage, not too much should be read into whether a species is described as being common, fairly common, uncommon, rare or very rare. It is never easy to assign descriptions such as these, and I readily admit that I have not adopted an entirely systematic approach in preparing this list! Also, the populations of many species will fluctuate from year to year or decade to decade.

This list should provide an important source of information for local moth recorders in helping to identify those species that are very rare and/or where records of such species might require confirmation. Obviously, other factors come into play, the recorder's experience and the potential for confusion with another species, but the main objective is to ensure that records of such species are validated as soon as possible to protect the long-term integrity of such records. There are also a number of particularly problematic species where we have far more unconfirmed records that confirmed records. These species have a symbol next to them. Please contact me by email or telephone if you think you have caught one of these species or any other species that might be considered unlikely or that has not been reliably recorded from the county before.

Unfortunately, this list may contain errors and omissions - please feel free to draw these to my attention.

Adrian Russell adrian@wainscot.demon.co.uk 0116 2415101

Ader Kolen Arzonica bicuspis April Kolen Arzonica da nin metototica Andre Molt Arzonica da nin metototica Andre Molt Ceraptery grammins Andre Molt Ceraptery	Alder Kitter Euroule historie	Days 9 Jacob
Angle Shades Phiogophora metaulosea  Range and mainly restricted to Charmwood Forest & actieme NW of County, Annual Charistia a beautratis  Annual Charistia a beautratis  Annual Charistia a beautratis  Local y common, accountly, Few recent records.  Annual Charistia a beautratis  Local y common, posteroid from Longshor in 2005.  Angent & Sable Rearmaphera healatta  August Thom Ennomes querienismis  Auturnia Charistia Charistia Industria Indus		
Angle-stepped Sallow Engarga paleaces Amuler Charitas obscurata Rare and restricted to western hard for county. Few recent records Anter Min Ceraptery, grammins Anter Min Ceraptery, grammins Anter Min Ceraptery, grammins Anter San Cerapter vestigation And Pug Eughheda innotata if risainata Auturnal More Perman auturnate Auturnal More Perman auturnate Auturnal More Perman auturnate Statu uncertain and anterior status of the status of t		
Annuel Kohrissa obscurata Annuel Kohrissa obscurata Antele Moth Craptery gramins Locally common particularly in Charavod Freez Livery more and Agreet Sable Poleracy agramins Locally common particularly in Charavod Freez Livery in 2005. Angues I Der Agrock substitution and the Charavod Sable Sa		
Anter Moth Corapteryx grammins Ancher's Darf Angolis evestigoidis Rev. VOH - 10 dipugnates Nicholar Gerords. Recorded from Loughtor in 2005. Angent & Sable Rheumaptere Instaltat Endroll - 12 ast record: Buddon Wood in 1962. Angent & Sable Rheumaptere Instaltat Endroll - 12 ast record: Buddon Wood in 1962. Angent & Sable Rheumaptere Instaltat Endroll - 12 ast record: Buddon Wood in 1962. Angent & Sable Rheumaptere Instaltat Endroll - 12 ast record: Buddon Wood in 1962. Angent & Sable Rheumaptere Instaltat Endroll - 12 ast record: Buddon Wood in 1962. Auturnal Angent Pertina auturnates  Auturnal Angent Pertina auturnates  Auturnal Rivest Euponorana giareosa Locally common in Charmacod Forest. Rare or absent elsewhere  Bannas Shawaya Antichioris enginis  Auturnal Auturnal Angent Pertina auturnates  Burred Cheshut Diasias dahili Very trac & declining. A Charmwood resident but only one post 1999 record.  Burred Cheshut Diasias dahili Very trac & declining. A Charmwood resident but only one post 1999 record.  Burred Cheshut Diasias dahili Very trac & declining. A Charmwood resident but only one post 1999 record.  Burred Workstonesis culturate  Uncommon.  Burred Workstonesis culturates  Uncommon.  Burred Workstonesis culturates  Uncommon.  Burred Workstonesis culturates  Common  Burred Workstonesis culturates  Common  Burred Workstonesis culturates  Common  Burred Workstonesis and trace as the service of the service and trace as the service as the service and trace as the servic	<u> </u>	
Acpert & Saber Remoration sessinguillas Anni Purguell' Establish Remoration in 1925. Anthorning Moht Epiritia subumbità Authurnal Rusbis Eugenorisma glarencesa Barnara Stowaway Antichioris eriphia Barred Chestrol Danisi darbiil Very rare & deciming. A Chamwood France Belearing. Barred Chestrol Danisi darbiil Very rare & deciming. A Chamwood France Belearing. Barred Chestrol Danisi darbiil Very rare & deciming. A Chamwood France Belearing. Barred Chestrol Danisi darbiil Very rare & deciming. A Chamwood France Belearing. Barred Chestrol Danisi darbiil Very rare & deciming. A Chamwood France Belearing. Danisi Chestrol Danisi darbiil Very rare & deciming. A Chamwood France Belearing. Danisi Chestrol Danisi darbiil Very rare & deciming. A Chamwood France Belearing. Danisi Chestrol Danisi darbiil Very rare & deciming. A Chamwood France Belearing. Danisi Chestrol Danisi darbiil Very rare & deciming. A Chamwood France Belearing. Danisi Chestrol Danisi darbiil Very rare & deciming. A Chamwood France Belearing. Danisi Chestrol Danisi darbiil Uncommon.  Barred Thoriti Champel Mercal Danisi darbiil Danisi Chestrol Danisi Chestrol Danisi darbiil Danisi Chestrol Danisi Chestrol Danisi Chestrol Danisi darbiil Danisi Chestrol Danis		
Angent & Sable Rheumaptera hastata An Pug Eupfliche innotata in Trainaha August Thom Ennomos queronaris August Thom Ennomos queronaris August Thom Ennomos queronaris August Thom Ennomos queronaris Beaufful Wey few records anno VCH, none in last 50 years.  Authurn Green Cappel Chinocolysis mistis Ennomos queronaris Beaufful Wey few records anno VCH, none in last 50 years.  Beaufful Wey few records anno VCH, none in last 50 years.  Beaufful Wey few records anno VCH, none in last 50 years.  Authurnal Rustic Eugenstain glarosa Barnaris Stowawy Articificitis grippina Advertives species. One record second on bannaise, 1985 Barred Chestrud Darsia dahili Barred Chestrud Darsia dahili Vey race & declaring, a Charmwood resident but only one post 1999 record.  Barred Real Phylase facciant Queronal States Chestrud Darsia dahili Vey race & declaring, a Charmwood resident but only one post 1999 record.  Barred States Blacka States and Chestral Common new Dramstel increase since mid 1970s, but declaring now?  Barred States Blacka States and post participation of the post post post post post post post post		
August Thom Ennomos quercinaria Auturn Green Capter Chiprocysts mists Enterd Tvey few seconds sane VEV, none in last 50 years. Auturnal Moht Epirita autumnata Auturnal Rust Euprociansi garreasa Sistaus uncertain, all records perhaps need investigation/verification Auturnal Rust Euprociansi garreasa Sistaus uncertain, all records perhaps need investigation/verification Auturnal Rust Euprociansi garreasa Auturnal Rust Euprociansi garreasa Sistaus uncertain, all records perhaps need investigation/verification Auturnal Rust Euprociansi garreasa Barread Fotole Parizona bilitacida Uncommon on Directain Garreasa Euprociansi Garreas	Argent & Sable Rheumaptera hastata	
Autumal Notice perital autumala  Saltau uncatari, all records partings need investigation/verification Autumal Nusic Eugonórima glareesa  Locally common in Charmwood Porest. Rare or absent elsewhere.  Barnara Stowway, Anfichoris enphisa Adventiva species. One record: se cocoon in banaras, 1986 Barned Chesthric Dansia dahili  Very rare 8 declining, A Charmwood resident but only one poster.  Barned Roth Pulyas specialis Barned Roth Pulyas specialis Barned Roth Pulyas specialis Barned Roth Pulyas specialis Barned Rowal Prestroma bifaciats Barned Rowal Prestroma bifaciats Barned Saltow Tillacea aurago Fairly common Barned Straw Ganderitis praviata Barned Very level peritoris Barned Villagedis pulyariata Barned Verlow Cidaria futuriata Barned Verlow Typacis pulyariata Barned Verlow Typacis	Ash Pug Eupithecia innotata f. fraxinata	
Auturneal Moth Epiritia autumnata Auturneal Moth Epiritia autumnata Auturneal Rustic Eugenorisma giareosa Locally common in Charmwood Forest Rare or absent elsewhere Banana Stowaway Antechnier enpha Aoventive speces. One record: ex occoor on bananas. 1985 Barred Cheshut Datasia darini Very sara & Destinga, Charmwood resident but only one post-1999 record. Director Mouldy Malescale under the Common & Localised, but now expanding the the northwest of the country. Barred Stowaway and the Perizona bifaciata Barred Stallow Titacea aurego Barred Stallow Titacea aurego Barred Stallow Titacea aurego Barred Stallow Titacea aurego Barred Stallow Gardents byralata Common Barred Uniber Plagodis puteraria Rared Stallow Gardent subvista Common Barred Uniber Plagodis puteraria Rared Stallow Gardent subvista Common Barred Window Calaria bulvata Common Beaudid Cheshut Agrochola pichnicia Beaudid Coleshut Agrochola pichnicia Beaudid Goden V Autographa pulchrina Beaudid Soden V Autographa pulchrina Beaudid Shout Hypena crassalis Beaudid Shout Hypena crassalis Very rare. Only four post-VCH records. Charmwood Lodge only colony? Beaudid Shout Hypena crassalis Very rare migrant. Last record in 2013. Beaudid Shout Hypena crassalis Very rare migrant. Last record in 2013. Beaudid Shout Hypena crassalis Very rare migrant. Last record in 2013. Beaudid Shout Hypena crassalis Very rare migrant. Last record in 2013. Beaudid Very last many last the subvision of the subvision		
Autumna Rustic Eugnorisma glarecsa Banrana Stowway Antichtors enphisa Alventives posese. One record co coopon on bananas, 1985 Barrad Chesthut Darisal dahil Very rare & declining, A. Charmwood resident but only one post-1999 record. Barred Horder, Washonalia culturaria Uncommon & Localised, but now expanding into the northwest of the county. Barred Ref Hystes (ascisant) Barred Salver Place (ascisant) Barred Salver Place (ascisant) Barred Salver Place (ascisant) Barred Salver Place (ascisant) Barred Vellow Cidara (buyta) Barred Hystes) Barred Hystes (buyta) Barred Vellow Cidara (buyta) Barred Hystes) Barred Hystes (buyta) Barred Hystes (buyta) Barred Vellow Cidara (buyta) Barred Hystes) Barred Hystes (buyta) Barred Hystes) Barred Hystes (buyta) Barred Hystes) Barred Hystes (buyta) Barred Hystes (buyta) Barred Hystes) Barred Hystes (buyta)		
Banana Stowaway Antichlors erphila Barred Chestrut Diaria dahili Very raza & declining, a Charmwood resident but only one post-1999 record. Barred Hock-lip Watsonalia culturaia Uncommon & Localised, but now expanding into the northwest of the county. Barred Red Hyber Basesiana Common now Dramatic increase since mid-1970's, but declining now? Barred Revinet Pertzona blassata Uncommon. Barred Revinet Pertzona blassata Uncommon. Barred Revinet Pertzona blassata Uncommon. Barred Warier Carda fulvata Common now Carda fulvata Barred Warier Carda fulvata Common Beaded Chestrut Agrochola lychnidis Barred Vellow Carda fulvata Common Beaded Chestrut Agrochola lychnidis Beaufful Carpet Messieuze alticiliata Uncommon and local in woodland. Beaufful Carpet Messieuze alticiliata Uncommon and local in woodland. Beaufful Hook-sp Laspeyria flexula Fairy common Fairy common Beaufful Hook-sp Laspeyria flexula Fairy common Beaufful Hook-sp Laspeyria flexula Fairy common Fairy common Beaufful Hook-sp Laspeyria flexula Fairy common Fairy common Beaufful Hook-sp Laspeyria flexula Fairy common Fairy common Fairy common Beaufful Hook-sp Laspeyria flexula Fairy common in west of Courty, Rare in Rutland & east. Black Rutland Laspes Agroched Laspes Spain deellista Fairy common in west of Courty, Rare in Rutland & east. Black Rutland Laspes Spain deellista Fairy common in Rutland woods, now becoming established in the west. Black Rutland Laspes Spain deellista Fairy common in Rutland woods, now becoming established in the west. Black Arches Lyvandria passible deellista Very rare migrant. Card flexular flexular flexu		
Barred Fock-thut Diarsia dahili  Very rare & deciling, A Charmwood resident but only one poet-1999 record.  Barred Red Hylase fasciaria  Locamon Now Dramstic increase since mid-1970's, but declining now?  Barred Red Hylase fasciaria  Common now. Dramstic increases since mid-1970's, but declining now?  Barred Salew Titaccea surgep  Faily common  Barred Salew Titaccea surgep  Barred Salew Titaccea surgep  Barred Salew Titaccea surgep  Faily common  Barred Salew Candents pyralsta.  Common  Barred Salew Cidaria fluidati  Common  Barred Salew Cidaria fluidati  Barred Salew Cidaria fluidati  Common  Beautiful Brocade Lacancible configua  Beautiful Goden Y Autographa pudchrina  Basulful Snoul Hypana crassalis  Very rare. Only four post-VCH records. Charmwood Lodge only colony?  Basulful Snoul Hypana crassalis  Very rare. Only four post-VCH records. Charmwood Lodge only colony?  Basulful Snoul Hypana crassalis  Very rare. Only four post-VCH records. Charmwood Lodge only colony?  Basulful Snoul Hypana crassalis  Very rare only four post-VCH records. Charmwood Lodge only colony?  Basulful Snoul Hypana crassalis  Very rare migrant. Last record in 2003.  Biotelyn-Pup Pashila debiata  Very rare migrant. Last record in 2004.  Biotelyn-Pup Pashila debiata  Very rare migrant. Last record in 2004.  Biotelyn-Pup Pashila debiata  Very rare migrant. Last record in 2005.  Biack Pashila debiata  Very rare migrant. Last record in 2005.  Biack Pashila debiata  Very rare migrant. Last record in 2006.  Common In Fluttend woods, now becoming established in the west, Now fairly common, naving recordly expanded its rarge county-wide  Biack Roules Lopedpula nigra  Now fairly common, naving recordly expanded its rarge county-wide  Biack Roules Lopedpula nigra sulphila debiata  Very rare migrant fo		
Barred Red Hyber Seasonian (Common & Durandic Increase since mid-1970's, but declining now?  Barred Red Nuiter Perizona bifosiata (Common on Durandic Increase since mid-1970's, but declining now?  Barred Salow Tilance aurago Fairly common (Common on Common	<u> </u>	
Barred Rivule Ferizoma Diffaciata Barred Sallow Tiliacea aurago Barred Sallow Tiliacea aurago Barred Sallow Tiliacea aurago Barred Sallow General Rivule Ferizoma Diffaciata Barred Sallow General Rivule Sallow Tiliacea aurago Barred Sallow General Rivule Sallow Sallow Sallow Tiliacea aurago Barred Oliver Plagodia pulverana Rare & malniy confined to Ulverscroft valley woodland. Common Barred Oliver Claria Infvata Barred Tollow Claria Infvata Barred Tollow Claria Infvata Barred Tollow Claria Infvata Beaufild Browde Laconolisa contiqua Beaufild Browde Laconolisa contiqua Beaufild Browde Laconolisa contiqua Beaufild Browde Laconolisa Contiqua Beaufild Golden Y Autographa pulchrina Beaufild South Hypena crassalis Beaufild South Hypena crassalis Beaufild South Hypena Crassalis Beaufild South Hypena Crassalis Very rare Most recent records are from Rutland. Beaufild South Hypena Crassalis Very rare Most recent records are from Rutland. Beaufild South Hypena Crassalis Very rare Most recent records are from Rutland. Beaufild South Hypena Crassalis Very rare Migrant. Last record in 2003. Beaufild South Hypena Crassalis Very rare migrant. Last record in 2003. Bid South William South		
Barred Stave Straw Gandarits preliated a Common Sarred Straw Straw Gandarits preliated Common Sarred Straw Gandarits preliated Common Sarred Straw Straw Gandarits preliated Common Garred There Plagode pulveraira Common Garred There Plagode Garred There Plagode Garred		Uncommon & localised, but now expanding into the northwest of the county.
Barred Staw Candartis pyraliata Barred Univer Plagodis pulveraria Barred Chestrut Agrochola lychnidie Beaufful Carpet Mesoleuca albicilista Beaufful Carpet Mesoleuca albicilista Beaufful Carpet Mesoleuca albicilista Beaufful Foot-gib Laspevia fiexulia Beaufful Hook-gib Laspevia fiexulia Beaufful Hook-gib Laspevia fiexulia Beaufful Hook-gib Laspevia fiexulia Beaufful Hook-gib Laspevia fiexulia Beaufful Fellow Underwrig Abarta mytilii Beaufful Fellow Beaufful Beaufful Fellow Beaufful Beaufful Fellow Beaufful Beaufful Fellow Beaufful Beaufful Beaufful Fellow Beaufful Beaufful Fellow Beaufful Beaufful Fellow Beaufful Beaufful Beaufful Fellow Beaufful Beaufful Beaufful Beaufful Fellow	,	
Barred Dimber Plagodis pulveraria Barred Vellow Cidaria fuhvata Barred Vellow Cidaria fuhvata Barred Vellow Cidaria fuhvata Barred Vellow Cidaria fuhvata Common Beaufful Agrochola lychndis Beaufful Agrochola lychndis Common Beaufful Brocade Leaenobia contigua Beaufful Golden P Autographa pulchrina Beaufful Snout hypena crassalis Very rare. Only four post-VCH records. Charmwood Lodge only colony? Beaufful Snout hypena crassalis Very rare. Only four post-VCH records. Charmwood Lodge only colony? Beaufful Snout hypena frama mythlii Very rare migrant. Last record in 2003. Beaufful Snout hypena frama mythlii Very rare migrant. Last record in 2003. Bulbery Pup Baphia debiliata Bulbery Pup Baphia de		
Barred Vellow Cidaria fulvata Bared Vellow Cidaria fulvata Bared Vellow Cidaria fulvata Baedaed Chestnut Agrochola lychnidis Common Baedaed Chestnut Agrochola lychnidis Baeufful Carpet Mesoleuca albicillata Baeufful Vellow Underwing Anafa myttili Baeufful Yellow Despetial Performance Bilbaeufful Yellow Underwing Anafa Myttili Baeufful Yellow Underwing Anafa Wallow Baeufful Yellow Underwing Anafa Wallow Baeufful Yellow Baeufful Yello		·
Barded Cheshrut Agrochola kyndridis  Peauliful Forcade Lacanobia contigua  Peauliful Forcade Lacanobia contigua  Peauliful Forcade Lacanobia contigua  Very rare. Most recent records are from Rutland.  Beauliful Golden Y Autographa pulchrina  Fairly common  Beauliful Golden Y Autographa pulchrina  Fairly common  Beauliful Snout Hyperna crassalia  Very rare. Only four post-VCH records. Charnwood Lodge only colony?  Rare. Only known from a single site near Ashty de la Zouch  Beauliful Snout Hyperna crassalia  Very rare migrant. Last record in 2003.  Biblierry Pup Fasphila debiletla  Bird's Wing Dypterygia scabriuscula  Bird's Morta Oxigonhora puppilaria  Uncommon in Wing wing recordity expanded fits range county-wide  Bird's Morta Oxigonhora puppilaria  Uncommon  Uncommon  Uncommon  Bird's Morta Oxigonhora puppilaria  Bird's Morta Oxigonhora puppilaria  Bird's Morta Oxigonhora puppilaria  Uncommon  Bird's Morta Oxigonhora puppilaria  Uncommon  Bird's Morta Oxigonhora puppilaria  Uncommon  Bird's Morta Oxigonhora puppilaria  Bird's Agra Oxigonhora puppilaria  Uncommon, Mainyl Yound in old woodland.  Uncommon Morta Oxigonhora puppilaria  Bord-ered Beauly Epione repandaria  Uncommon, Mainyl Yound in damp habitats. Occasional in gardens.  Bord-ered Pupp Euphreia abrovata  Bord-ered Siraw Heliothia publig		
Beautiful Foxed Leannabla contiqua  Beautiful Carget Mesoleuza albicillata  Beautiful Carget Mesoleuza albicillata  Uncommon and local in woodland.  Beautiful Hook-tip Laspeyris flexula  Beautiful Foxed typespyris flexula  Beatiful Foxed typespyris flexula  Locally common in single site near Ashby de la Zouch  Very rare Only known from a single site near Ashby de la Zouch  Very rare Only known from a single site near Ashby de la Zouch  Very rare Only foxed typespyris flexula  Locally common in general Capter Flexula  Locally common in Beatiful Known from a single site near Ashby de la Zouch  Very rare Only foxed typespyrish flexula  Locally common in Beatiful Known from a		
Beautiful Grocarde Lacanobia contigua  Peautiful Golden Y Autographa putchrina  Beautiful Golden Y Autographa putchrina  Beautiful Golden Y Autographa putchrina  Beautiful Snout Hypena crassalis  Very rare. Only four post-VCH records. Charwood Lodge only colony?  Beautiful Snout Hypena crassalis  Very rare. Only four post-VCH records. Charwood Lodge only colony?  Beautiful Snout Hypena crassalis  Very rare. Only four post-VCH records. Charwood Lodge only colony?  Beautiful Snout Hypena crassalis  Very rare migrant. Last record in 2003.  Biblitery Pup Pasphila debiliata  Extinc? Last record. Sienfield, 1939.  Bird's Wing Dypterygia scabriuscula  Biack Arches Lymantria monacha  Locally common in Rutland woods, now becoming established in the west.  Black Rustic Aportophyra ingra  Black Arches Lymantria monacha  Locally common in Rutland woods, now becoming established in the west.  Black Rustic Aportophyra ingra  Black Arches Lymantria monacha  Locally common in Rutland woods, now becoming established in the west.  Black Rustic Aportophyra ingra  Now fairly common, having recently expended its range county-wide  Blackneck Lyeppla pastimum  Uncommon.  Blair's Shoulder-ton Lithipohane leautieri  Very rare migrant. Only one record, from an Edith Weston garden in 2006.  Blackneck Lyeppla pastimum  Very rare sitatus uncertain. All records need investigating.  Blood-ven't Timandra comae  Common in gardens.  Common in gardens.  Blood-ven't Timandra comae  Common in gardens.  Rare a Note of the stream of the stre		
Beautiful Carpet Mesoleuca albiciliata Beautiful Fook-lip Laspeyria flexula Beautiful Fook-lip Laspeyria flexula Beautiful Fook-lip Laspeyria flexula Beautiful Yellow Underwing Anarta myrtilli Beautiful Yellow Underwing Anarta myrtilli Beautiful Yellow Underwing Anarta myrtilli Beatiful Yellow Underwing Anarta myrtilli Biberry Pug Pasiphila debiliata Biberry Pug Pasiphila debiliata Bird's Wing Dyptergia scaphiuscula Black Arches Lymantria monacha Locally common in Rutland woods, now becoming established in the west. Black Arches Lymantria monacha Locally common in Rutland woods, now becoming established in the west. Black Rustic Appropriya in a common in west of County, Rare in Rutland & east. Black Rustic Appropriya in a common in Rutland woods, now becoming established in the west. Black Rustic Appropriya in a common in service of County, Rare in Rutland & east. Discommon in Rutland woods, now becoming established in the west. Black Rustic Appropriya in a common in service of County, Rare in Rutland & east. Discommon in Rutland woods, now becoming established in the west. Black Rustic Appropriya in a construction of the service of County, Rare in Rutland & east. Discommon in Rutland woods, now becoming established in the west. Discommon in Rutland woods, now becoming established in the west. Discommon in Rutland woods, now becoming established in the west. Discommon in Rutland woods, now becoming established in the west. Discommon in Rutland woods, now becoming established in the west. Discommon in Rutland woods, now becoming established in the west. Discommon in Rutland woods, now becoming established in the west. Discommon in Rutland woods, now becoming established in the west. Discommon in Rutland woods, now becoming established in the west. Discommon in gardens. Discommon in Rutland woods, now becoming established in the west. Discommon in Rutland woods, now becoming established in the west. Discommon in Rutland woods, now becoming established in the west. Discommon in Rutland woods, now becoming establis		
Beautiful Golden Y Autorgrapha pulchrina  Fairly common  Beautiful Snout Hypena crassalis  Very rare. Only four post-VCH records. Charwood Lodge only colony?  Beautiful Snout Hypena crassalis  Very rare. Only four post-VCH records. Charwood Lodge only colony?  Beautiful Yellow Underwing Ananta myntili  Bedstraw Hawk-moth Hybes galii  Very rare migrant. Last record in 2003.  Bilbierry Pup Pasphila debilata  Extinct? Last record. Selnefield, 1939.  Bird's Wing Dypterygia scabriuscula  Black Arches Lymantria monacha  Locally common in west of County, Rare in Rutland & east.  Black Rustic Aporophyla nigra  Black Arches Lymantria monacha  Locally common in west of County, Rare in Rutland & east.  Black Rustic Aporophyla nigra  Now fairty common in west of County, Rare in Rutland & east.  Locally common in Wild woods, now becoming established in the west.  Locally common in Wild woods, now becoming established in the west.  Locally common in Burland woods, now becoming established in the west.  Locally common in Burland woods, now becoming established in the west.  Locally common in Burland woods, now becoming established in the west.  Locally common in Burland woods, now becoming established in the west.  Locally common in Burland woods, now becoming established in the west.  Locally common in Burland woods, now becoming established in the west.  Locally common in Burland woods, now becoming established in the west.  Locally common in Burland woods, now becoming established in the west.  Locally common in Burland woods, now becoming established in the west.  Locally common in Burland woods, now becoming established in the west.  Locally common in Burland woods, now becoming established in the west.  Locally common in Burland woods, now becoming established in the west.  Locally common in Burland woods and an analysis of the provided woods and an an analysis of the provided woods and an an an analysis of the provided woods and an an an analysis of the provided woods and an an an analysis of the provided woods and an		
Beautiful Hook tip Laspeyria flexula  Fairly common.  Beautiful Yellow Underwing Anarta mytilli  Beatiful Yellow Underwing Anarta mytilli  Bedstraw Hawk-moth Hyles galli  Very rare migrant. Last record in 2003.  Bilberny Pug Pasiphila debilata  Extinct? Last record: Glenfield, 1939.  Bilack Arches Lynantria monacha  Black Arches Lynantria monacha  Black Arches Lynantria monacha  Black Arches Lynantria monacha  Black Rustle Apoprophyla nigra  Black Rustle Apoprophyla nigra  Nov fairly common in Rutland woods, now becoming established in the west.  Black Rustle Apoprophyla nigra  Nov fairly common in Rutland woods, now becoming established in the west.  Black Rustle Apoprophyla nigra  Nov fairly common in Rutland woods, now becoming established in the west.  Black Rustle Apoprophyla nigra  Nov fairly common in Rutland woods, now becoming established in the west.  Black Rustle Apoprophyla nigra  Nov fairly common in Rutland woods, now becoming established in the west.  Black Rustle Apoprophyla nigra  Nov fairly common in Rutland woods, now becoming established in the west.  Black Rustle Apoprophyla nigra in Nov fairly common in Rutland woods, now becoming established in the west.  Black Rustle Apoprophyla nigra in Nov fairly common in Rutland woods, now becoming established in the west.  Black Rustle Apoprophyla nigra in Nov fairly common in Rutland woods, now becoming established in the west.  Black Rustle Apoprophyla nigra in Nov fairly common in Rutland woods, now becoming established in the west.  Black Rustle Apoprophyla nigra in Rustle Rustl		
Beautiful Snout Hypena crassalis  Bedstaw Hawk-moth Hybes galili  Extinct? Last record in 2003.  Birds Wing Dypterygia scabriuscula  Biack Rushic Aporophyla nigra  Biack Arches Lymantria monacha  Locally common in west of County, Rare in Rutland & east.  Biack Arches Lymantria monacha  Locally common in Mulland woods, now becoming established in the west.  Now fairly common, having recently expanded its range county-wide  Biackneck Lygephila pastinum  Uncommon.  Biair's Mocha Cyclophora puppillaria  Biar's Bhouder-knot Lithophan leautieri  Common in gardens.  Bieached Pug Eupitheoia expalidata  Very rare migrant. Only one record, from an Edith Weston garden in 2006.  Biesched Pug Eupitheoia expalidata  Very rarestistus uncertain. All records need investigating.  Biodos-wen Timandra comae  Common  Biossom Underwing Orthosia miniosa  Biossom Underwing Statis uncertain.  Biard Rama di Campaton Diagnamia  Biossom Underwing Orthosia miniosa  Broadered Pug Eupitheoia Succentratia  Broadered Pug Eupitheoia  Broadered Pug Eupi		Fairly common.
Beautiful Yeliow Underwing Anarta myrtilli Bedsfraw Hawk-moth Hyles galli Very rare migrant. Last record in 2003.  Bilberny Pug Pasiphila debilata Bird's Wing Dyteryiga scabriuscula Bilack Arches Lymantira monacha Bilack Arches Lymantira monacha Bilack Arches Lymantira monacha Bilack Rustlo. Aporophyla nigra Bilack Sylophila pastinum Uncommon. Bilar's Shouder-knot Lithophane leaulteri Bilar's More and the strain leaulteri Bilar's More and leaulteri Bilar's More and localised primarily found in damp habitats. Ocasional in gardens. Bordered Stale uplineder alucentural		Very rare. Only four post-VCH records. Charnwood Lodge only colony?
Bedstraw Hawk-moth Hyles galili Bilbern Pup Pasphilal debilata Bird's Wing Dypterygia scabriuscula Biack Arches Lymantria monacha Locally common in Wast of County, Rare in Rutland & east. Biack Rustic Aporophyla nigra Biack Arches Lymantria monacha Locally common in Wast of County, Rare in Rutland & east. Biack Rustic Aporophyla nigra Biack Arches Lymantria monacha Locally common in Wast of County, Rare in Rutland & east. Biack Rustic Aporophyla nigra Biack Arches Lymantria monacha Locally common in Rustland woods, now becoming established in the west. Now fairly common, having recently expanded its range county-wide Biaris Shoulder-knot Lithophan leautier Common. Biaris Shoulder-knot Lithophan leautier Common in gardens. Biack Arches Lathan and Carbiacha and Ca		Rare. Only known from a single site near Ashby de la Zouch
Bird's Wing Dypterygia scabriuscula Black Arches Lymartria monacha Locally common in Ritland woods, now becoming established in the west. Black Rustic Aporophyla nigra Black Arches Lymartria monacha Locally common in Ritland woods, now becoming established in the west. Now fairly common, having recently expanded its range county-wide Blackneck Lypephila pastirum Blair's Mocha Cyclophora puppillaria Blair's Shoulder-knot Lithophane leautieri Bleached Pug Eupithecia expaliidata Very rare sitratus uncertain. All records need investigating. Common in gardens. Bleached Pug Eupithecia expaliidata Very raresistatus uncertain. All records need investigating. Common Blood-wein Timandra comae Brood-wein Timandra Common Brood-wein Timandra comae Brood-wein Timandra Common Brood-bordered Bee Hawk-moth Hemaris fuoliomae Brood-bordered Vellow Underweing Noctua timbrata Brown-barred Carpet Electrophaes corylata B		Very rare migrant. Last record in 2003.
Black Arches Lymantria monacha Black Rustic Aporophyla nigra Black Rustic Aporophyla nigra Black Rustic Aporophyla nigra Black Rustic Aporophyla nigra Black Rustic Acyclophora puspiliaria Blair's Mocha Cyclophora puspiliaria Blair's Mocha Cyclophora puspiliaria Blair's Mocha Cyclophora puspiliaria Blair's Mocha Cyclophora puspiliaria Black Rustic Acyclophora puspiliaria Black Rustic Acyclophora puspiliaria Black Rustic Architecture and State		,
Blackneck Lygephila pastinum  Blar's Mocha Cyclophora puppillaria  Blar's Mocha Cyclophora puppillaria  Blar's Mocha Cyclophora puppillaria  Blar's Mocha Cyclophora puppillaria  Very rare migrant. Only one record, from an Edith Weston garden in 2006.  Blar's Shouder-knot Lithophane leautieri  Common in gardens.  Bleached Pug Eupithecia expallidata  Very rare/status uncertain. All records need investigating.  Blood-vein Timardra comae  Blossom Underwing Orthosia miniosa  Blotched Emerald Comibaena bajularia  Uncommon, maniny in damp habitats  Drodered Beauty Epione repandaria  Uncommon, Malniy found in damp habitats. Occasional in gardens.  Bordered Pug Eupithecia succenturiata  Brordered Pug Eupithecia succenturiata  Brordered Straw Hellottis peltigera  Drodered White Bupatus primaria  Locally common, but possibly declining in recent years.  Bright-line Brown-eye Lacanobia oleracea  Brinstone Moth Opisthograptis luteolata  Brindled Green Dryobotodes eremita  Brindled White-spot Paradarisa extersaria  Broad-bordered White Hecatera bicciorata  Broad-bordered White Hecatera bicciorata  Broad-bordered White Hecatera bicciorata  Broad-bordered Vellow Underwing Moctua Imbrista al Rocken-barred Carpet Electrophaes corylata  Brown Music Rusina ferrugina  Brown Music Rusina ferrugina  Brown Music Rusina ferrugina  Brown Stellophalia Rusina ferrugina  Brown St		
Blackneck Lygephila pastinum    Uncommon.		
Blair's Mocha Cyclophora puppillaria Blair's Shoulder-knot Lithophane leaulieri Common in gardens. Bleached Pug Eupithecia expallidata Very rarer/status uncertain. All records need investigating. Blood-vein Timandra comae Blood-vein Timandra comae Blood-vein Timandra comae Blossom Underwing Orthosia miniosa Rare. VCH and recent records from Rutland and Dadlington. Rare and localised: primarily found in old woodland. Blotched Emerald Combaena bajularia Blue-bordered Carpet Plemyria rubiginata Bordered Beauty Epione repandaria Bordered Beauty Epione repandaria Bordered Beauty Epione repandaria Bordered Beauty Epione repandaria Bordered Pug Eupithecia succenturiata Bordered Pug Eupithecia succenturiata Bordered Pug Eupithecia succenturiata Bordered Straw Heliothis petitigera Bordered Straw Heliothis petitigera Uncommon, but possibly declining in recent years. Brick Agrochola circellaris Brick Agrochola circellaris Bright-line Brown-eye Lacanobia oleracea Brinstone Moth Opisthograptis luteolata Common Brindled Gene Dryobotodes eremita Brindled Other Dasypojia tempil Brindled Other Dasypojia tempil Brindled White-spot Paradarisa extersaria Broad-bordered Bee Hawk-moth Hemaris fuciformis Broad-bordered Yellow Underwing Noctua fimbriata Broad-bordered Yellow Underwing Noctua fimbriata Brown-spot Plantera Brindled Green Carpet Electrophaes corylata Brown-spot Philora Republica Somplia Prown-spot Brindled Green Carpet Electrophaes corylata Brown-spot Philora Republica Somplia Prown-spot Philora Republication Rep	1 1 2 5	
Blairs Shoulder-knot Lithophane leautier  Bleached Pug Eupithecia expedilidata  Very rare/status uncertain. All records need investigating.  Blood-vein Timandra comae  Blossom Underwing Orthosia miniosa  Blotched Emerald Comibaena bajularia  Blotched Emerald Comibaena bajularia  Blotched Emerald Comibaena bajularia  Blotched Emerald Comibaena bajularia  Blot-ordered Carpet Piemyria rubiginata  Uncommon, found mainly in damp habitats  Bordered Garpet Piemyria rubiginata  Uncommon, found mainly found in damp habitats  Bordered Gothic Sidendis reticulata  Bordered Gothic Sidendis reticulata  Bordered Sallow Pyrnia umbra  Bordered Sallow Pyrnia umbra  Bordered Sallow Pyrnia umbra  Bordered White Bupalus piniaria  Birck Agrochola circellaris  Bright-line Brown-eye Lacanobia oleracea  Brimstone Molth Opisthograpits luteolata  Brindled Green Dryobotodes eremita  Brindled Green Dryobotodes eremita  Brindled Green Dryobotodes eremita  Brindled Green Dryobotodes eremita  Brindled Pug Eupithecia subreviata  Broad-bordered Bee Hawk-moth Hemaris fuciformis  Broad-bordered Bee Hawk-moth Hemaris fuciformis  Broad-bordered Bee Hawk-moth Hemaris fuciformis  Brown-Brown Brown Hemaris fuciformis  Brown-Brown Brown Bro		
Bleached Pug Eupithecia expallidata  Very rare/status uncertain. All records need investigating.  Common  Blossom Underwing Orthosia miniosa  Blotched Emeratd Comibaena bajularia  Rare and localised: primarily found in old wooddand.  Blue-bordered Carpet Plemyria rubiginata  Bordered Beauty Epione repandaria  Bordered Beauty Epione repandaria  Bordered Beauty Epione repandaria  Bordered Pug Eupithecia succenturiata  Bordered Pug Eupithecia succenturiata  Bordered Pug Eupithecia succenturiata  Bordered Straw Heliothis peltigera  Bordered Straw Heliothis peltigera  Uncommon, but possibly declining  Bordered Straw Heliothis peltigera  Uncommon migrant.  Bordered Straw Heliothis peltigera  Uncommon migrant.  Locally common, but possibly declining in recent years.  Fairly common  Brindled Beauty Lycia Initaria  Common  Brindled Beauty Lycia Initaria  Brindled Green Dryobtodes eremita  Brindled Oren Dayspolia tempil  Findled Oren Dayspolia tempil  Findled White-spot Paradarisa extersaria  Broad-barder White Hecatera bioclorata  Broad-bordered White Hecatera bioclorata  Broad-bordered Bee Hawk-moth Hemaris fuciformis  Broad-bordered Stellow Underwing Noctua fimbriata  Brown-Brown Rustic Rusina ferruginea  Brown Sulley Delineria abbrovata  Brown Sulley Delineria position sulla a very rare. Only one crocer: Cuembrough garden (2008)  Brown Sulley Phileme vetulata  Brown Sulley Phileme vetulata  Brown-Sulley Phileme Persophora chlorosata  Frairly common in susteen half of County, uncommon in east.  Formerly common, particularly in Rutland. Less common lesswhere.  Formerly common, particularly in Rutland & east Leics.  Brown-Sulley Phileme Persophora chlorosata  Formerly Common, particularly in Gramwood Forest  Formerly Common in suitable habitats	7 1 11	
Blossom Underwing Orthosia miniosa Blossom Underwing Orthosia miniosa Blote-bet Emerald Comibaena bajularia Blue-bordered Carpet Plemyria rubiginata Uncommon, found mainly found in old woodland. Blue-bordered Carpet Plemyria rubiginata Uncommon, found mainly found in old woodland. Blue-bordered Gothic Sideridis reticulata Bordered Beauty Epione repandaria Uncommon, fladingh plabitats. Occasional in gardens. Bordered Gothic Sideridis reticulata Bordered Sallow Pyrnha umbra Bordered Sallow Pyrnha umbra Bordered Sallow Pyrnha umbra Bordered Sallow Pyrnha umbra Bordered Wara Heliothis petigera Uncommon migrant. Uncommon Brindled Green Dryobtodes eremita Brindled Pug Eupithecia abbreviata Brindled Pug Eupithecia abbreviata Common Brindled White-spot Paradarisa extersaria Broad-bordered Bee Hawk-moth Hemaris fuciformis Broad-bordered Bee Hawk-moth Hemaris fuciformis Broad-bordered Yellow Underwing Noctua fimbriata Broad-bordered Yellow Underwing Noctua fimbriata Brown Holla (Breamia pisis Broad-bordered Yellow Underwing Noctua fimbriata Brown Holla (Breamia pisis Brown Hemaris Fire prophora chitorosata Brown Scallop Philereme vetulata Uncommon in swatsem half of County, uncommon in east. Very rare. Only one record: Quentiborough garden (2008). Eraity common in swatsem half of County, uncommon in chara		
Blossom Underwing Orthosia miniosa Blotched Emerald Comibaena bajularia Rare and localised: primarily found in old woodland. Blue-bordered Carpet Plemyria rubiginata Uncommon, found mainly in damp habitats Bordered Beauty Epione repandaria Uncommon. Mainly found in in damp habitats. Occasional in gardens. Extinct? No records for 40+ years. Bordered Pug Eupithecia succenturata Bordered Pug Eupithecia succenturata Bordered Straw Heliothis peltigera Bordered Straw Heliothis peltigera Bordered White Bupalus prinata Brick Agrochola circellaria Fairly common, but possibly declining Bordered White Bupalus piniana Locally common, but possibly declining Brick Agrochola circellaria Fairly common Bright-line Brown-eye Lacanobia oleracea Brindshord Mor Opisthographis futerolata Common Brindled Beauty Lycia hirtaria Brindled Green Dryobotodes eremita Brindled Green Dryobotodes eremita Brindled Othre Dasypolia tempil Very rare. Recorded only once: two from Rutland Water (2004) Brindled Pug Eupithecia abbreviata Broad-bordered Bee Hawk-moth Hemaris fuciformis Broad-bordered Bee Hawk-moth Hemaris fuciformis Broom Moth Ceramica pisi Fairly common in gardens Broom Moth Ceramica pisi Fairly common in gardens Brown Rutland Strushia Fairly common Brindled White Rupalus piniana Broken-barred Carpet Electrophaes corylata Brown Rutland Strushia ferruginea Brown Scallop Philereme vetulata Common Brown Scallop Philereme vetulata Locally common, particularly in Rutland, Less common elsewhere. Brown-line Bright-eye Mythinna conigera Brown-line Bright-eye Mythinna conigera Brown-line Bright-eye Mythinna conigera Common in suitable habitats, particularly in Charnwood Forest Brown-line Bright-eye Mythinna conigera Common in suitable habitats. Brown-line Bright-eye Mythinna conigera Common in suitable habitats. Brown-line Bright-eye Mythinna conigera Uncommon but increased in last 10 years. Brown-line Bright-eye Mythinna conigera Uncommon but increased in last 10 years. Brown-line Bright-eye Mythinna conigera Uncommon but increased in last 10		
Blotched Emerald Comibaena bajularia Blue-bordered Carper Plempria rubiginata Uncommon, found mainly in damp habitats Bordered Beauty Epione repandaria Bordered Gohlie Sidendis reticulata Bordered Sollie Sidendis reticulata Bordered Sollie Sidendis reticulata Bordered Sug Eupithecia succenturiata Bordered Sallow Pyrhia umbra Bordered White Bupalus piniaria Brick Agrochoa circellaris Bright-line Brown-eye Lacanobia oleracea Common Brimslone Moth Opisthograpis lutelolata Common Brimslone Moth Opisthograpis lutelolata Common Brindled Green Dryobtodes eremita Brindled Green Dryobtodes eremita Brindled Pug Eupithecia abbreviata Brindled Pug Eupithecia abbreviata Brindled Pug Eupithecia abbreviata Broad-barred White Hecatera bicolorata Broad-bordered Bee Hawk-moth Hemaris fuciformis Broad-bordered Yellow Underwing Noctua fimbriata Brown-barred Carpet Electrophaes corylata Brown Moth Ceramica pisi Broom Moth Ceramica pisi Brown Moth Ceramica pisi Brown Moth Ceramica pisi Brown Silver-line Petrophora chiorosata Brown Silver-line Petrophora chiorosata Brown Silver-line Petrophora chiorosata Brown-lip Chesias rufata Very rare. Only one record: Queniborough garden (2008). Brown Silver-line Petrophora chiorosata Brown-lip Chesias rufata Very rare. Only one record: Queniborough garden (2008). Brown-lip Chesias rufata Very rare. Only one record: Queniborough garden (2008). Brown-lip Chesias rufata Very rare. Only one record: Queniborough garden (2008). Brown-lip Chesias rufata Very rare. Only one record: Queniborough garden (2008). Brown-lip Chesias rufata Very rare. Only one record: Queniborough garden (2008). Brown-lip Chesias rufata Very rare. Only one record: Queniborough garden (2008). Brown-lip Chesias rufata Very rare. Very rare. Only one record: Queniborough garden (2008). Brow		
Blue-bordered Carpet Plemyria rubiginata  Bordered Beauty Epione repandaria  Bordered Schick Sidendis reticulata  Bordered Gothic Sidendis reticulata  Bordered Pug Eupithecia succenturiata  Bordered Pug Eupithecia succenturiata  Bordered Straw Heliothis petitigera  Locally common, but possibly declining in recent years.  Brick Agrochola circellaris  Brick Agrochola circellaris  Brick Agrochola circellaris  Bright-line Brown-eye Lacanobia oleracea  Brindstone Moth Opisthograptis luteolata  Brindled Beauty Lycia hirtaria  Brindled Green Dryobotodes eremita  Brindled Ochre Dasypolia templi  Brindled Ochre Dasypolia templi  Frindled Pug Eupithecia abbreviata  Brindled Pug Eupithecia abbreviata  Broad-bordered Bee Hawk-moth Hemaris fuciformis  Broad-bordered Bee Hawk-moth Hemaris fuciformis  Broad-bordered Bee Hawk-moth Hemaris fuciformis  Brown House Rusina ferruginea  Brown Rusite Rusina ferruginea  Brown Rusite Rusina ferruginea  Brown Rusite Rusina ferruginea  Brown Scallop Philereme vetulata  Brown Scallop Philereme vetulata  Brown Scallop Philereme vetulata  Brown Scallop Philereme vetulata  Brown-line Bright-eye Mythimna conigera  Brown-spot Prinon Agrochola iltura  Formyn-spot Rinon Agrochola iltura  Formyn-veined Wainscot Archanara dissoluta  Brown-veined Wainscot Archanara dissoluta  Brown-spot Pinon Agrochola iltura  Formity common  Buff Formine Spilosoma lutea  Uncommon but increased in last 10 years.  Buff Ermine Spilosoma lutea  Uncommon but increased in last 10 years.  Buff Pelera bucephala  Buff Forma Euclidia glyphica  Uncommon negraticularly in Agrochola Nocot Rutland & east Leics.  Brown-spot Pinon Agrochola iltura  Formeriy common  Formity common  Buff Forma Putland & east Leics.  Brown-line Bright-eye Mythima conigera  Drommon Butland Water, but occurs elsewhere  Brussels Laco Cleorodes Bichenaria  Uncommon but in	U	
Bordered Beauty Epione repandaria Dordered Gohibc Sideridis reticulata Extinct? No records for 40+ years.  Bordered Gohibc Sideridis reticulata Bordered Sallow Pyrnha umbra Bordered Sallow Pyrnha umbra Bordered Sallow Pyrnha umbra Bordered Straw Heliothis petigera Duncommon migrant.  Bordered White Bupalus piniaria Locally common, but possibly declining Bordered White Bupalus piniaria Locally common, but possibly declining in recent years.  Brick Agrochola circellaris Brick Agrochola circellaris Bright-line Brown-eye Lacanobia oleracea Brinstone Moth Opisthograptis luteolata Common Brindled Green Dryobotodes eremita Brindled Green Dryobotodes eremita Brindled Green Dryobotodes eremita Brindled Ochre Dasypolia templi Very rare. Recorded only once: two from Rutland Water (2004) Brindled White-spot Paradarisa extersaria Broad-bordered White Hecatera bicolorata Broad-bordered Bee Hawk-moth Hemaris fuciformis Broad-bordered Bee Hawk-moth Hemaris fuciformis Brown-barred Carpet Electrophaes conylata Brown-barred Carpet Electrophaes conylata Broom-tip Chesias rufata Broom-tip Chesias rufata Broom-tip Chesias rufata Common Brown Scallop Philereme vertulata Brown Scallop Philereme vertulata Brown-Sizel Philora chiorosata Brown-Sizel Rusina ferruginea Brown-Brinnia Petrophora chiorosata Brown-Bright-eye Mythimna conigera Brown-Bright-eye Mythimna conigera Brown-Fizel Philereme vertulata Brown-Bright-eye Mythimna conigera Brown-Bright-eye Mythimna conigera Brown-Bright-eye Mythimna conigera Common Brown-Sallop Philereme vertulata Brown-spile Petrophora chiorosata Uncommon Mainly recorded from Rutland Water, but occurs elsewhere Brown-line Bright-eye Mythimna conigera UVCH an landful of recent records. Localised in Rutland & east Leics. Brown-spile Philerene vertulata Brow		
Bordered Gothic Sideridis reticulata Bordered Pug Eupithecia succenturiata Bordered Sallow Pyrrhia umbra Bordered Sallow Pyrrhia umbra Bordered Straw Heliothis peltigera Bordered White Bupalus piniaria Brick Agrochola circellaris Brick Agrochola circellaris Brick Agrochola circellaris Brish Locally common, but possibly declining in recent years.  Common Brindled Beauty Lycia hitratia Common Brindled Beauty Lycia hitratia Common Brindled Ochre Dasypolia templi Brindled Ochre Dasypolia templi Brindled Ochre Dasypolia templi Brindled White-spot Paradarisa extersaria Brindled White-spot Paradarisa extersaria Broad-bordered Bee Hawk-moth Hemaris fuciformis Broad-bordered Bee Hawk-moth Hemaris fuciformis Broad-bordered Vellow Underwing Noctua fimbriata Brown Hustic Rusina ferruginea Brown Rustic Rusina ferruginea Brown Rustic Rusina ferruginea Brown Rustic Rusina ferruginea Brown Rustic Rusina ferruginea Brown Scallop Philereme vetulata Brow		
Bordered Pug Eupithecia succenturiata Bordered Sallow Pyrrhia umbra Bordered Straw Heliothis peltigera Uncommon migrant. Locally common, but possibly declining in recent years. Brick Agrochola circellaris Bright-line Brown-eye Lacanobia oleracea Brimstone Moth Opisthographis luteolata Common Brindled Green Dryobotodes eremita Brindled Green Dryobotodes eremita Brindled Green Dryobotodes eremita Brindled Orber Dasypolia templi Very rare. Recorded only once: two from Rutland Water (2004) Brindled Pug Eupithecia abbreviata Brindled White-spot Paradarisa extersaria Broad-bordered Sele Hawk-moth Hemaris fuciformis Broad-bordered Vellow Underwing Noctua fimbriata Broad-bordered Vellow Underwing Noctua fimbriata Broom-tip Chesias rufata Broom-tip Chesias rufata Broom-tip Chesias rufata Broom-tip Chesias rufata Brown Scallop Philereme vetulata Brown Scallop Philereme vetulata Brown Scallop Philereme vetulata Brown Signly-line Petrophora chiorosata Brown-Signly-Philereme vetulata Brown-Birght-eye Mythimna conigera Brown-spilos phinon Agrochola litura Brown-reine Wainscot Archanara dissoluta Brown-reine Wainscot Archanara dissoluta Brown-reine Bright-eye Mythimna conigera Brown-spilos phinon Agrochola litura Brown-spilos p	, , ,	
Bordered Straw Heliothis peltigera Bordered White Bupalus piniaria Bordered White Bupalus piniaria Locally common, but possibly declining in recent years.  Fairly common Bringht-line Brown-eye Lacanobia oleracea Common Brindled Beauty Lycia hirtaria Brindled Green Dryobtodoes eremita Brindled Green Dryobtodoes eremita Brindled Ochre Dasypolia templi Very rare. Recorded only once: two from Rutland Water (2004) Brindled Pug Eupithecia abbreviata Common Brindled White-spot Paradarisa extersaria Broad-bordered Bee Hawk-moth Hemaris fuciformis Broad-bordered Bee Hawk-moth Hemaris fuciformis Broom Moth Ceramica pisi Broom Moth Ceramica pisi Brown Moth Ceramica pisi Brown Rustic Rusina ferruginea Brown Rustic Rusina ferruginea Brown Silver-line Petrophora chiorosata Brown-Spot Pinion Agrochola litura Brown-spot Pinion Agrochola litura Brown-spot Pinion Agrochola litura Brown-spot Pinion Agrochola litura Buff Arches Habrosyne pyritoides Buff-lish Phalera bucketha Fairly common Buff Footman Eliema depressa Burnished Brass Diadrhyspita Philiphale Burnished Brass Diadrhyspita Philiphale Burnished Brass Diadrhyspita Philiphale Burnished Bross Diadrhyspita Philiphale Burnished Bross Diadrhyspita Philiphale Burnished Brass Diadrhyspita Buff-tip Phalera bucephala Burnished Brass Diadrhyspita Buff-tip Phalera bucephala Burnished Brass Diadrhyspita Buff-tip Phalera bucephala Burnished Brass Diadrhysia Common Butterbur Hydraecia petasitis Parily common (Buddon Wood) & Ketton Quarry (2015) only		
Bordered White Bupalus piniaria Brick Agrochola circellaris Fairly common Fright-line Brown-eye Lacanobia oleracea Common Brinstone Moth Opisthograpits luteolata Common Brindled Green Dryobotodes eremita Brindled Green Dryobotodes eremita Brindled Green Dryobotodes eremita Brindled Ochre Dasypolia tempii Brindled Ochre Dasypolia tempii Brindled Ochre Dasypolia tempii Brindled Wilte-spot Paradarisa extersaria Brindled White-spot Paradarisa extersaria Brindled White-spot Paradarisa extersaria Broad-bordered Bee Hawk-moth Hemaris fuciformis Broad-bordered Bee Hawk-moth Hemaris fuciformis Broad-bordered Bee Hawk-moth Hemaris fuciformis Broom Moth Ceramica pisi Brown Rustic Rusina ferruginea Brown Rustic Rusina ferruginea Common Brown Rustic Rusina ferruginea Common Brown Scallop Philereme vetulata Brown-Silver-line Petrophora chorosata Brown-Ipi Chesias rufata Common Brown-Ipi Chesias rufata Common Brown-Bright-eye Mythimna conigera Common Brown-Pot Pinion Agrochola itura Brown-Pusi Petrophora diorasata Brown-Iali Euproctis chrysornioea Common is unitable habitats. Brown-Iali Euproctis ch		
Brick Agrochola circellaris Bright-line Brown-eye Lacanobia oleracea Common Brimstone Moth Opisthographis luteolata Common Brindled Beauty Lycia hirtaria Common Brindled Green Dryobotodes eremita Brindled Ochre Dasypolia templi Very rare. Recorded only once: two from Rutland Water (2004) Brindled Pug Eupithecia abbreviata Common Brindled White-spot Paradarisa extersaria Brindled White-spot Paradarisa extersaria Broad-bordered Bee Hawk-moth Hemaris fuciformis Broad-bordered Bee Hawk-moth Hemaris fuciformis Broad-bordered Bee Hawk-moth Hemaris fuciformis Broad-bordered Yellow Underwing Noctua fimbriata Broken-barred Carpet Electrophaes corylata Broken-barred Carpet Electrophaes corylata Brom-tip Chesias rufata Very rare. Only one record: Queniborough garden (2008). Brown Rustic Rusina ferruginea Common. Brown Scallop Philereme vetulata Locally common, particularly in Rutland. Less common elsewhere. Brown-line Bright-eye Mythimna conigera Brown-spot Pinion Agrochola litura Brown-spot	Bordered Straw Heliothis peltigera	Uncommon migrant.
Bright-line Brown-eye Lacanobia oleracea Common Brimstone Moth Opisthograptis luteolata Common Brindled Beauty Lycia hirtaria Common Brindled Green Dryobotodes eremita Brindled Green Dryobotodes eremita Brindled Ochre Dasypolia templi Brindled Que Eupithecia abbreviata Common Brindled White-spot Paradarisa extersaria Brindled White-spot Paradarisa extersaria Broad-barred White Hecatera bicolorata Broad-barred White Hecatera bicolorata Broad-bordered Bee Hawk-moth Hemaris fuciformis Broad-bordered Green Beeven Beeven Beeven Berind? Novemon in gardens Broad-bordered Green Beeven Beeven Berind? Novemon Brown-barred Carpet Electrophaes corylata Fairly common Brown Rustic Rusina ferruginea Common Brown Rustic Rusina ferruginea Common Brown Rustic Rusina ferruginea Common Brown-Bailop Philereme vetulata Locally common, particularly in Rutland, Less common elsewhere. Brown-Both Pinion Agrochola litura Formerly common, particularly in Rutland, Less common elsewhere.  Brown-spot Pinion Agrochola litura Formerly common, now only fairly common  Brown-tail Euproctis chrysorrhoea VCH & a handful of recent records. Localised in Rutland & east Leics.  Brown-barred Carpen Bervesa Uncommon but increased in last 10 years.  Buff Arches Habrosyne pyritoides Common Buff Emine Spilosoma lutea Uncommon but increased in last 10 years.  Buff-lip Phalera bucephala Fairly	Bordered White Bupalus piniaria	Locally common, but possibly declining in recent years.
Brimstone Moth Opisthograptis luteolata Common Brindled Beauty Lycia hirtaria Common Brindled Green Dryobotodes eremita Brindled Cohre Dasypolia templi Very rare. Recorded only once: two from Rutland Water (2004) Brindled Pug Eupithecia abbreviata Brindled White-spot Paradarisa extersaria Rare. Only recorded from Burbage Wood area and a few Rutland sites. Broad-barred White Hecatera bicolorata Broad-bordered Bee Hawk-moth Hemaris fuciformis Broad-bordered Yellow Underwing Noctua fimbriata Broad-bordered Yellow Underwing Noctua fimbriata Broad-bordered Yellow Underwing Noctua fimbriata Broken-barred Carpet Electrophaes corylata Broom-tip Chesias rufata Very rare. Only one record: Queniborough garden (2008). Brown Rustic Rusina ferruginea Brown Scallop Philerem vetulata Brown Scallop Philerem vetulata Brown-line Bright-eye Mythimna conigera Brown-spot Phinon Agrochola litura Brown-spot Pinion Agrochola litura Brown-spot Pinion Agrochola litura Brown-spot Pinion Agrochola litura Brown-veined Wainscot Archanara dissoluta Uncommon. Brussels Lace Cleorodes lichenaria Buff Arches Habrosyne pyritoides Buff Ermine Spilosoma lutea Buff Arches Habrosyne pyritoides Uncommon Buff Ermine Spilosoma lutea Buff-tip Phalera bucephala Burnet Companion Euclidia glyphica Burnets Arches Habrosyne pyritoides Buff-tip Phalera bucephala Burnet Companion Euclidia glyphica Butterbur Hydraecia petasitis Extinct? NCH secorded for Manareas. Fairly common Fairly common Fairly common Localised in last 10 years. Fairly common Butterbur Hydraecia petasitis Extinct? VCH & a couple of other records. Under-recorded? Very rare. Only one records. Fairly common Fairly common Butterbur Hydraecia petasitis Extinct? VCH & a couple of other records. Under-recorded? Very rare. Only one records. Fairly common Fairly common Fairly common Fairly common Fairly common in suitable habitats Fairly common Uncommon/under-recorded. Most frequently recorded from Rutland & east. Fairly common Fairly common Fairly common Fairly common Fairly common Fairly common Fairl		
Brindled Green Dryobotodes eremita Brindled Ochre Dasypolia templi Very rare. Recorded only once: two from Rutland Water (2004)  Brindled Pug Eupithecia abbreviata Brindled White Hecatera bicolorata Broad-barred White Hecatera bicolorata Broad-barred White Hecatera bicolorata Broad-bordered Bee Hawk-moth Hemaris fuciformis Broad-bordered Pellow Underwing Noctua fimbriata Broad-bordered Yellow Underwing Noctua fimbriata Broad-bordered Yellow Underwing Noctua fimbriata Broad-bordered Yellow Underwing Noctua fimbriata Brown-barred Carpet Electrophaes corylata Brown Moth Ceramica pisi Brown Moth Ceramica pisi Brown Scallop Philereme vetulata Brown Scallop Philereme vetulata Brown Scallop Philereme vetulata Brown-Silver-line Petrophora chlorosata Brown-spot Pinion Agrochola litura Brown-spot Pinion Agrochola litura Brown-veined Wainscot Archanara dissoluta Brussels Lace Cleorodes lichenaria Uncommon. Buff Arches Habrosyne pyritoides Buff Fairly common Buff Frairle Carpet Selectrophaes Common Buff Frairle Carpet Selectrophaes Common Buff-phalera bucephala Buff-phalera bucephala Burnet Companion Euclidia glyphica Burnished Brass Diachrysia chrystits Common Butterbur Hydraecia petasitis Extinct? No recorded from Burbage Wood area and a few Rutland sites. Brairly common in garricy common in gears thou substance and a few Rutland sites. Brairly common Fairly common Fairly common  Brairly common in western half of County, uncommon in east. Very came on western half of County, uncommon in east. Very rare. Only one record: Queniborough garden (2008). Brairly common in western half of County, uncommon in east. Very rare. Only one record: Queniborough garden (2008). Brairly common in suitable habitats, particularly in Rutland. Less common elsewhere. Very common in suitable habitats, particularly in Charmwood Forest Brown-spot Pinion Agrochola litura Formerly common, now only fairly common Brown-tail Euproctis chrysorrhoea VCH & a handful of recent records. Localised in Rutland & east Leics. Brown-spot Pinion Agrochola litura		
Brindled Green Dryobotodes eremita Brindled Ochre Dasypolia templi Brindled Ochre Dasypolia templi Brindled Pug Eupithecia abbreviata Common Brindled White-spot Paradarisa extersaria Broad-barred White Hecatera bicolorata Broad-bordered Bee Hawk-moth Hemaris fuciformis Broad-bordered Yellow Underwing Noctua fimbriata Broad-bordered Yellow Underwing Noctua fimbriata Broad-bordered Yellow Underwing Noctua fimbriata Broom-barred Carpet Electrophaes corylata Broom-bordered Searnica pisi Broom-bip Chesias rufata Brown-bip Chesias rufata Brown Rustic Rusina ferruginea Brown Scallop Philereme vetulata Brown-line Bright-eye Mythimna conigera Brown-spot Pinion Agrochola litura Brown-tail Euproctis chrysormhoea Brown-tail Euproctis chrysormhoea Brown-tail Euproctis chrysormhoea Brown-tail Euproctis chrysormhoea Buff Arches Habrosyne pyritoides Buff Fraining Phalera bucephala Buff Fraining Agrochola litura Buff Fraining Phalera bucephala Burnish Mainscot Nonagria typhae Buttoned Snout Hypena rostralis Extinct? No recorded from Burbage Wood area and a few Rutland sites. Brown-line Bright-eye Mythimna conigera Common in gardens Fairity common Brown-tail Euproctis chrysormhoea Very rare. Only one record: Queniborough garden (2008). Brown-line Bright-eye Mythimna conigera Common in suitable habitats, particularly in Charnwood Forest Brown-tail Euproctis chrysormhoea Very common in suitable habitats. Brown-eine Wainscot Archanara dissoluta Uncommon. Mainly recorded from Rutland & east Leics. Brown-eined Wainscot Archanara dissoluta Uncommon but increased in last 10 years. Buff-tip Phalera bucephala Fairly common Bulrush Wainscot Nonagria typhae Uncommon/under-recorded. Most frequently recorded from Rutland & east. Burnished Brass Diachrysia chrysitis Extinct? VCH & a couple of other records. Under-recorded? Buttoned Snout Hypena rostralis Extinct? VCH & a couple of other records. Under-recorded? Buttoned Snout Hypena rostralis Very rare. VCH record (Buddon Wood) & Ketton Quarry (2015) only		
Brindled Ochre Dasypolia templi Very rare. Recorded only once: two from Rutland Water (2004) Brindled Pug Eupithecia abbreviata Common Brindled Multie-spot Paradarisa extersaria Rare. Only recorded from Burbage Wood area and a few Rutland sites. Broad-barred White Hecatera bicolorata Fairly common in gardens Broad-bordered Bee Hawk-moth Hemaris fuciformis Extinct? No records for nearly 20 years & those may be doubtful. Broad-bordered Yellow Underwing Noctua fimbriata Fairly common. Broad-bordered Yellow Underwing Noctua fimbriata Fairly common. Broken-barred Carpet Electrophaes corylata Fairly common Broom Moth Ceramica pisi Fairly common in western half of County, uncommon in east. Broom-tip Chesias rufata Very rare. Only one record: Queniborough garden (2008). Brown Rustic Rusina ferruginea Common. Brown Scallop Philereme vetulata Locally common, particularly in Rutland. Less common elsewhere. Brown Silver-line Petrophora chlorosata Very common in suitable habitats, particularly in Charnwood Forest Brown-lse Bright-eye Mythimna conigera Common in suitable habitats, particularly in Charnwood Forest Brown-tail Euproctis chrysorrhoea VCH & a handful of recent records. Localised in Rutland & east Leics. Brown-veined Wainscot Archanara dissoluta Uncommon. Mainly recorded from Rutland Water, but occurs elsewhere WCH only. Buff Arches Habrosyne pyritoides Common Buff Footman Eilema depressa Uncommon but increased in last 10 years. Buff-tip Phalera bucephala Fairly common Bultrush Wainscot Nonagria typhae Common, especially in damp habitats Burnet Companion Euclidia glyphica Uncommon/under-recorded. Most frequently recorded from Rutland & east. Brown-lead Brass Diachrysia chrysitis Extinct? VCH & a couple of other records. Under-recorded? Buttoned Snout Hypena rostralis Very rare. VCH record (Buddon Wood) & Ketton Quarry (2015) only Cabbage Moth Mamestra brassicae		
Brindled Pug Eupithecia abbreviata Brindled White-spot Paradarisa extersaria Broad-barred White Hecatera bicolorata Broad-bordered Bee Hawk-moth Hemaris fuciformis Broad-bordered Bee Hawk-moth Hemaris fuciformis Broad-bordered Pellow Underwing Noctua fimbriata Broken-barred Carpet Electrophaes corylata Broom Moth Ceramica pisi Broom-tip Chesias rufata Broom-tip Chesias rufata Brown-tip Chesias rufata Brown Rustic Rusina ferruginea Brown Scallop Philereme vetulata Brown-line Bright-eye Mythimna conigera Brown-spot Pinion Agrochola litura Brown-spot Pinion Agrochola litura Brown-veined Wainscot Archanara dissoluta Brown-veined Wainscot Archanara dissoluta Buff Arches Habrosyne pyritoides Buff Footman Ellema depressa Buff-tip Phalera bucephala Bulrush Wainscot Nonagria typhae Bulrush Wainscot Nonagria typhae Buttender Nonagria typnae Buttenbur Hydraecia petasitis Very rare. VCH compon, pericularly in Rutland Waeter, but occurs elsewhere Common Butterbur Hydraecia petasitis Very rare. Very ecorded from Rutland Water, but occurs elsewhere Brusses Lace Cleorodes lichenaria VCH only. Buff Arches Habrosyne pyritoides Common Buff Emmine Spilosoma lutea Common Buff Footman Ellema depressa Uncommon but increased in last 10 years. Buff-tip Phalera bucephala Fairly common Buff Emmon Fairly common Fairly common Buff Fairly common Fairly common Buff Fairly common Fairly for the recorded Most frequently recorded from Rutland & east. Burnet Companion Euclidia glyphica Uncommon Fairly Common Fairly Common Buff Fairly common Fairly Common Fairly Common Buff Fairly Common Fairly		
Brindled White-spot Paradarisa extersaria Broad-barred White Hecatera bicolorata Broad-bordered Bee Hawk-moth Hemaris fuciformis Broad-bordered Yellow Underwing Noctua fimbriata Broken-barred Carpet Electrophaes corylata Broom Moth Ceramica pisi Broom-tip Chesias rufata Brown-tip Chesias rufata Brown Sallop Philereme vetulata Brown Silver-line Petrophora chlorosata Brown-line Bright-eye Mythimna conigera Brown-tail Euproctis chrysorrhoea Brown-veined Wainscot Archanara dissoluta Bruff Fortman Eilema depressa Buff-fip Phalera bucephala Buff-fip Phalera bucephala Buff-fip Palera bucephala Buttoned Snout Hypena rostralis Common Butterbur Hydraecia petasitis Extinct? No recorded from Burbage Wood area and a few Rutland stes. Fairly common in gardens Extinct? No records for nearly 20 years & those may be doubtful.  Extinct? No records for nearly 20 years & those may be doubtful.  Extinct? No records for nearly 20 years & those may be doubtful.  Fairly common.  Fairly common  Fairly common  Fairly common in western half of County, uncommon in east.  Fairly common in western half of County, uncommon in east.  Fairly common in western half of County, uncommon in east.  Fairly common.  Fairly common.  Fairly common in western half of County, uncommon in east.  Fairly common.  Scallop Phalera bucephala Fairly common.  Fairly common in western half of County, uncommon elsewhere.  Very rare. Only one record: Queniborough garden (2008).  Fairly common.  Fairly common Buttern half of County, uncommon in east.  Fairly common.  Fairly common.  Fairly common in western half of County, uncommon in east.  Fairly common.  Fairly common.  Fairly common in western half of County, uncommon in east.  Fairly common.  Fairly common.  Fairly common.  Fairly common in western half of County, uncommon in east.  Fairly common.		
Broad-barred White Hecatera bicolorata Broad-bordered Bee Hawk-moth Hemaris fuciformis Broad-bordered Yellow Underwing Noctua fimbriata Broken-barred Carpet Electrophaes corylata Broom Moth Ceramica pisi Broom-tip Chesias rufata Brown-tip Chesias rufata Brown Scallop Philereme vetulata Brown Scallop Philereme vetulata Brown-line Bright-eye Mythimna conigera Brown-spot Pinion Agrochola litura Brown-veined Wainscot Archanara dissoluta Bruff Fortman Eilema depressa Buff-tip Phalera bucephala Burnet Companion Euclidia glyphica Buttoned Snout Hypena rostralis Buttoned Snout Hypena rostralis Broad-bordered Bee Hawk-moth Hemaris fuciformis Extinct? No records for nearly 20 years & those may be doubtful. Extinct? No records for nearly 20 years & those may be doubtful. Extinct? No records for nearly 20 years & those may be doubtful. Extinct? No records for nearly 20 years & those may be doubtful. Extinct? No records for nearly 20 years & those may be doubtful. Extinct? No records for nearly 20 years & those may be doubtful. Extinct? No records for nearly 20 years & those may be doubtful.  Extinct? No records for nearly 20 years & those may be doubtful.  Extinct? No records for nearly 20 years & those may be doubtful.  Extinct? No records for nearly 20 years & those may be doubtful.  Extinct? No records for nearly 20 years & those may be doubtful.  Extinct? No records for nearly 20 years & those may be doubtful.  Extinct? No records for nearly 20 years & those may be doubtful.  Extinct? No records for nearly 20 years & those may be doubtful.  Extinct? No records for nearly 20 years & those may be doubtful.  Extinct? No records for nearly 20 years & those may be doubtful.  Extinct? No records for nearly 20 years & those may be doubtful.  Extinct? No records for nearly 20 years & those may be doubtful.  Extinct? No records for nearly 20 years & those may be doubtful.  Extinct? No records for nearly 20 years & those may be doubtful.  Extinct Parly common in susteden habitats and particularly in Rutland & east Lecs.  Uncommo		
Broad-bordered Bee Hawk-moth Hemaris fuciformis Broad-bordered Yellow Underwing Noctua fimbriata Broken-barred Carpet Electrophaes corylata Fairly common.  Fairly common.  Fairly common in western half of County, uncommon in east.  Broom-tip Chesias rufata Very rare. Only one record: Queniborough garden (2008).  Brown Rustic Rusina ferruginea Common.  Brown Scallop Philereme vetulata Locally common, particularly in Rutland. Less common elsewhere.  Brown Silver-line Petrophora chlorosata Very common in suitable habitats, particularly in Charnwood Forest Common in suitable habitats.  Brown-spot Pinion Agrochola litura Brown-tail Euproctis chrysorrhoea VCH & a handful of recent records. Localised in Rutland & east Leics.  Brown-veined Wainscot Archanara dissoluta Uncommon. Mainly recorded from Rutland Water, but occurs elsewhere VCH only.  Buff Arches Habrosyne pyritoides Common Buff Footman Eilema depressa Uncommon but increased in last 10 years.  Buff-tip Phalera bucephala Buff-tip Phalera bucephala Suff-tip Phalera bucephala Common Buff Sourned Rutland Sourhysits Common Common Euclidia glyphica Uncommon/under-recorded. Most frequently recorded from Rutland & east.  Burnet Companion Euclidia glyphica Uncommon/under-recorded. Most frequently recorded?  Very common Very common Very common Very common, nespecially in urban areas.		,
Broad-bordered Yellow Underwing Noctua fimbriata Broken-barred Carpet Electrophaes corylata Fairly common Fairly common Fairly common in western half of County, uncommon in east. Fairly common in western half of County, uncommon in east. Fairly common in western half of County, uncommon in east. Fairly common in western half of County, uncommon in east. Forom-tip Chesias rufata Very rare. Only one record: Queniborough garden (2008).  Brown Rustic Rusina ferruginea Common. Frown Scallop Philereme vetulata Locally common, particularly in Rutland. Less common elsewhere. Forom Silver-line Petrophora chlorosata Forom-line Bright-eye Mythimna conigera Common in suitable habitats. Formerly common, now only fairly common Frown-tail Euproctis chrysorrhoea Forom-veined Wainscot Archanara dissoluta Formerly common, now only fairly common Frown-veined Wainscot Archanara dissoluta Uncommon. Mainly recorded from Rutland & east Leics.  WCH only.  Buff Arches Habrosyne pyritoides Common Buff Fortman Eilema depressa Uncommon but increased in last 10 years.  Buff-tip Phalera bucephala Fairly common, especially in damp habitats  Burnet Companion Euclidia glyphica Uncommon/under-recorded. Most frequently recorded from Rutland & east.  Burnished Brass Diachrysia chrysitis Common Butterbur Hydraecia petasitis Extinct? VCH & a couple of other records. Under-recorded?  Extinct? VCH & a couple of other records. Under-recorded?  Fairly common, especially in urban areas.		
Broken-barred Carpet Electrophaes corylata Broom Moth Ceramica pisi Broom Moth Ceramica pisi Broom-tip Chesias rufata Broom-tip Chesias rufata Broom-tip Chesias rufata Brown Rustic Rusina ferruginea Brown Rustic Rusina ferruginea Brown Scallop Philereme vetulata Brown Silver-line Petrophora chlorosata Brown-spot Pinion Agrochola litura Brown-spot Pinion Agrochola litura Brown-tail Euproctis chrysorrhoea Brown-eined Wainscot Archanara dissoluta Brussels Lace Cleorodes lichenaria Buff Arches Habrosyne pyritoides Buff Forman Eilema depressa Buff-tip Phalera bucephala Buff-tip Phalera bucephala Burnet Companion Euclidia glyphica Butterbur Hydraecia petasitis Butterbur Hydraecia petasitis Very rare. Only one record: Queniborout, uncommon in east. Very rare. Only one record: Queniborout, uncommon in east. Very rare. Only one record: Queniborout, uncommon in east. Very rare. Only one record: Queniborout, uncommon in east. Very rare. Only one record: Queniborout, uncommon in east. Very rare. Only one record: Queniborout, uncommon elsewhere (2008). Buff avily common in suitable habitats, particularly in Charnwood Forest Common, now only fairly common  Uncommon. Mainly recorded from Rutland & east Leics. Uncommon but increased in Rutland Water, but occurs elsewhere VCH only. Cabbage Moth Mamestra brassicae  Fairly common  Uncommon but increased in last 10 years.  Butterout Gudon Wood) & Ketton Quarry (2015) only Cabbage Moth Mamestra brassicae		
Broom Moth Ceramica pisi Fairly common in western half of County, uncommon in east. Broom-tip Chesias rufata Very rare. Only one record: Queniborough garden (2008). Brown Rustic Rusina ferruginea Common. Brown Scallop Philereme vetulata Locally common, particularly in Rutland. Less common elsewhere. Very common in suitable habitats, particularly in Charnwood Forest Brown-line Bright-eye Mythimna conigera Common in suitable habitats. Brown-spot Pinion Agrochola litura Formerly common, now only fairly common Brown-tail Euproctis chrysorrhoea VCH & a handful of recent records. Localised in Rutland & east Leics. Brown-veined Wainscot Archanara dissoluta Uncommon. Mainly recorded from Rutland Water, but occurs elsewhere VCH only. Buff Arches Habrosyne pyritoides Common Buff Ermine Spilosoma lutea Common Buff Footman Eilema depressa Uncommon but increased in last 10 years. Buff-tip Phalera bucephala Fairly common Bulrush Wainscot Nonagria typhae Common, especially in damp habitats Burnet Companion Euclidia glyphica Uncommon/under-recorded. Most frequently recorded from Rutland & east. Common Butterbur Hydraecia petasitis Extinct? VCH & a couple of other records. Under-recorded? Buttoned Snout Hypena rostralis Very rare. VCH record (Buddon Wood) & Ketton Quarry (2015) only Cabbage Moth Mamestra brassicae	<u>_</u>	
Brown-tip Chesias rufata Very rare. Only one record: Queniborough garden (2008). Brown Rustic Rusina ferruginea Common. Brown Scallop Philereme vetulata Locally common, particularly in Rutland. Less common elsewhere. Brown Silver-line Petrophora chlorosata Very common in suitable habitats, particularly in Charnwood Forest Common in suitable habitats. Brown-line Bright-eye Mythimna conigera Common in suitable habitats. Brown-spot Pinion Agrochola litura Formerly common, now only fairly common Brown-tail Euproctis chrysorrhoea VCH & a handful of recent records. Localised in Rutland & east Leics. Brown-veined Wainscot Archanara dissoluta Uncommon. Mainly recorded from Rutland Water, but occurs elsewhere VCH only. Buff Arches Habrosyne pyritoides Common Buff Froitman Eilema depressa Uncommon but increased in last 10 years. Buff-tip Phalera bucephala Fairly common Bulrush Wainscot Nonagria typhae Common, especially in damp habitats Burnet Companion Euclidia glyphica Uncommon/under-recorded. Most frequently recorded from Rutland & east. Burnished Brass Diachrysia chrysitis Common Butterbur Hydraecia petasitis Extinct? VCH & a couple of other records. Under-recorded? Very rare. VCH record (Buddon Wood) & Ketton Quarry (2015) only Cabbage Moth Mamestra brassicae		
Brown Scallop Philereme vetulata  Brown Silver-line Petrophora chlorosata  Brown-line Bright-eye Mythimna conigera  Brown-spot Pinion Agrochola litura  Brown-tail Euproctis chrysorrhoea  Brown-veined Wainscot Archanara dissoluta  Brunseis Lace Cleorodes lichenaria  Buff Footman Eilema depressa  Buff-tip Phalera bucephala  Burnet Companion Euclidia glyphica  Burnshed Brass Diachrysia chrysitis  Butterbur Hydraecia petasitis  Brown-spot Pinion Agrochola litura  Common  Formerly common, now only fairly common  VCH & a handful of recent records. Localised in Rutland & east Leics.  Uncommon. Mainly recorded from Rutland Water, but occurs elsewhere  VCH only.  Common  Buff Ermine Spilosoma lutea  Common  Buff Footman Eilema depressa  Uncommon but increased in last 10 years.  Buff-tip Phalera bucephala  Fairly common  Bulrosh Wainscot Nonagria typhae  Common, especially in damp habitats  Uncommon/under-recorded. Most frequently recorded from Rutland & east.  Burnished Brass Diachrysia chrysitis  Butterbur Hydraecia petasitis  Extinct? VCH & a couple of other records. Under-recorded?  Very rare. VCH record (Buddon Wood) & Ketton Quarry (2015) only  Cabbage Moth Mamestra brassicae		
Brown Scallop Philereme vetulata  Brown Silver-line Petrophora chlorosata  Brown-line Bright-eye Mythimna conigera  Brown-spot Pinion Agrochola litura  Brown-tail Euproctis chrysorrhoea  Brown-veined Wainscot Archanara dissoluta  Brunseis Lace Cleorodes lichenaria  Buff Footman Eilema depressa  Buff-tip Phalera bucephala  Burnet Companion Euclidia glyphica  Burnshed Brass Diachrysia chrysitis  Butterbur Hydraecia petasitis  Brown-spot Pinion Agrochola litura  Common  Formerly common, now only fairly common  VCH & a handful of recent records. Localised in Rutland & east Leics.  Uncommon. Mainly recorded from Rutland Water, but occurs elsewhere  VCH only.  Common  Buff Ermine Spilosoma lutea  Common  Buff Footman Eilema depressa  Uncommon but increased in last 10 years.  Buff-tip Phalera bucephala  Fairly common  Bulrosh Wainscot Nonagria typhae  Common, especially in damp habitats  Uncommon/under-recorded. Most frequently recorded from Rutland & east.  Burnished Brass Diachrysia chrysitis  Butterbur Hydraecia petasitis  Extinct? VCH & a couple of other records. Under-recorded?  Very rare. VCH record (Buddon Wood) & Ketton Quarry (2015) only  Cabbage Moth Mamestra brassicae	Brown Rustic Rusina ferruginea	
Brown-line Bright-eye Mythimna conigera  Brown-spot Pinion Agrochola litura  Brown-spot Pinion Agrochola litura  Brown-tail Euproctis chrysorrhoea  VCH & a handful of recent records. Localised in Rutland & east Leics.  Brown-veined Wainscot Archanara dissoluta  Uncommon. Mainly recorded from Rutland Water, but occurs elsewhere  Brussels Lace Cleorodes lichenaria  VCH only.  Buff Arches Habrosyne pyritoides  Buff Ermine Spilosoma lutea  Common  Buff Footman Eilema depressa  Uncommon but increased in last 10 years.  Buff-tip Phalera bucephala  Bulrush Wainscot Nonagria typhae  Burnet Companion Euclidia glyphica  Burnished Brass Diachrysia chrysitis  Burnished Brass Diachrysia chrysitis  Butterbur Hydraecia petasitis  Extinct? VCH & a couple of other records. Under-recorded?  Buttoned Snout Hypena rostralis  Very rare. VCH record (Buddon Wood) & Ketton Quarry (2015) only  Cabbage Moth Mamestra brassicae  Fairly common, especially in urban areas.	Brown Scallop Philereme vetulata	
Brown-spot Pinion Agrochola litura  Brown-tail Euproctis chrysorrhoea  Brown-veined Wainscot Archanara dissoluta  Brussels Lace Cleorodes lichenaria  Buff Arches Habrosyne pyritoides  Buff Footman Eilema depressa  Buff-tip Phalera bucephala  Bulrush Wainscot Nonagria typhae  Burnet Companion Euclidia glyphica  Burnished Brass Diachrysia chrysitis  Butterbur Hydraecia petasitis  Buttoned Snout Hypena rostralis  Formerly common, now only fairly common  VCH & a handful of recent records. Localised in Rutland & east Leics.  Uncommon. Mainly recorded from Rutland Water, but occurs elsewhere  VCH only.  Common  Common  Uncommon but increased in last 10 years.  Fairly common  Common, especially in damp habitats  Uncommon/under-recorded. Most frequently recorded from Rutland & east.  Common  Butterbur Hydraecia petasitis  Extinct? VCH & a couple of other records. Under-recorded?  Buttoned Snout Hypena rostralis  Very rare. VCH record (Buddon Wood) & Ketton Quarry (2015) only  Cabbage Moth Mamestra brassicae  Fairly common, especially in urban areas.		
Brown-tail Euproctis chrysorrhoea  Brown-veined Wainscot Archanara dissoluta  Brussels Lace Cleorodes lichenaria  Buff Arches Habrosyne pyritoides  Buff Footman Eilema depressa  Buff-tip Phalera bucephala  Burnet Companion Euclidia glyphica  Burnished Brass Diachrysia chrysitis  Butterbur Hydraecia petasitis  Brown-veined Wainscot Nonagria typea  VCH only.  Common  Common  Uncommon but increased in last 10 years.  Fairly common  Common, especially in damp habitats  Uncommon/under-recorded. Most frequently recorded from Rutland & east.  Common  Butterbur Hydraecia petasitis  Extinct? VCH & a couple of other records. Under-recorded?  Buttoned Snout Hypena rostralis  VCH only.  Common  Common  Buff Footman Eilema depressa  Uncommon but increased in last 10 years.  Fairly common  Common, especially in damp habitats  Uncommon/under-recorded. Most frequently recorded from Rutland & east.  Common  Butterbur Hydraecia petasitis  Extinct? VCH & a couple of other records. Under-recorded?  Buttoned Snout Hypena rostralis  Very rare. VCH record (Buddon Wood) & Ketton Quarry (2015) only  Cabbage Moth Mamestra brassicae  Fairly common, especially in urban areas.		
Brown-veined Wainscot Archanara dissoluta  Brussels Lace Cleorodes lichenaria  Buff Arches Habrosyne pyritoides  Buff Footman Eilema depressa  Buff-tip Phalera bucephala  Burnet Companion Euclidia glyphica  Burnished Brass Diachrysia chrysitis  Butterbur Hydraecia petasitis  Buttoned Snout Hypena rostralis  VCH only.  Common  Common  Uncommon but increased in last 10 years.  Fairly common  Common, especially in damp habitats  Uncommon/under-recorded. Most frequently recorded from Rutland & east.  Common  Extinct? VCH & a couple of other records. Under-recorded?  Buttoned Snout Hypena rostralis  Very rare. VCH record (Buddon Wood) & Ketton Quarry (2015) only  Cabbage Moth Mamestra brassicae		
Brussels Lace Cleorodes lichenaria  WCH only.  Buff Arches Habrosyne pyritoides  Buff Ermine Spilosoma lutea  Buff Footman Eilema depressa  Buff-tip Phalera bucephala  Bulrush Wainscot Nonagria typhae  Burnet Companion Euclidia glyphica  Burnished Brass Diachrysia chrysitis  Butterbur Hydraecia petasitis  Buttoned Snout Hypena rostralis  Common, especially in damp habitats  Uncommon/under-recorded. Most frequently recorded from Rutland & east.  Common  Butterbur Hydraecia petasitis  Extinct? VCH & a couple of other records. Under-recorded?  Buttoned Snout Hypena rostralis  Very rare. VCH record (Buddon Wood) & Ketton Quarry (2015) only  Cabbage Moth Mamestra brassicae  Fairly common, especially in urban areas.		
Buff Arches Habrosyne pyritoides  Buff Ermine Spilosoma lutea  Buff Footman Eilema depressa  Buff-tip Phalera bucephala  Bulrush Wainscot Nonagria typhae  Burnet Companion Euclidia glyphica  Burnished Brass Diachrysia chrysitis  Butterbur Hydraecia petasitis  Buttoned Snout Hypena rostralis  Common, especially in damp habitats  Uncommon/under-recorded. Most frequently recorded from Rutland & east.  Common  Extinct? VCH & a couple of other records. Under-recorded?  Buttoned Snout Hypena rostralis  Very rare. VCH record (Buddon Wood) & Ketton Quarry (2015) only  Cabbage Moth Mamestra brassicae  Common, especially in urban areas.		
Buff Ermine Spilosoma lutea  Buff Footman Eilema depressa  Buff-tip Phalera bucephala  Bulrush Wainscot Nonagria typhae  Burnet Companion Euclidia glyphica  Burnished Brass Diachrysia chrysitis  Butterbur Hydraecia petasitis  Buttoned Snout Hypena rostralis  Common, especially in damp habitats  Uncommon/under-recorded. Most frequently recorded from Rutland & east.  Common  Extinct? VCH & a couple of other records. Under-recorded?  Buttoned Snout Hypena rostralis  Very rare. VCH record (Buddon Wood) & Ketton Quarry (2015) only  Cabbage Moth Mamestra brassicae  Fairly common, especially in urban areas.		,
Buff Footman Eilema depressa  Buff-tip Phalera bucephala  Bulrush Wainscot Nonagria typhae  Burnet Companion Euclidia glyphica  Burnished Brass Diachrysia chrysitis  Butterbur Hydraecia petasitis  Buttoned Snout Hypena rostralis  Cabbage Moth Mamestra brassicae  Uncommon but increased in last 10 years.  Fairly common  Common, especially in damp habitats  Uncommon/under-recorded. Most frequently recorded from Rutland & east.  Common  Extinct? VCH & a couple of other records. Under-recorded?  Very rare. VCH record (Buddon Wood) & Ketton Quarry (2015) only  Cabbage Moth Mamestra brassicae  Fairly common, especially in urban areas.		
Buff-tip Phalera bucephala  Bulrush Wainscot Nonagria typhae  Burnet Companion Euclidia glyphica  Burnished Brass Diachrysia chrysitis  Butterbur Hydraecia petasitis  Buttoned Snout Hypena rostralis  Cabbage Moth Mamestra brassicae  Fairly common  Fairly common  Common  Lucommon/under-recorded. Most frequently recorded from Rutland & east.  Common  Extinct? VCH & a couple of other records. Under-recorded?  Very rare. VCH record (Buddon Wood) & Ketton Quarry (2015) only  Cabbage Moth Mamestra brassicae  Fairly common, especially in urban areas.		
Bulrush Wainscot Nonagria typhae Common, especially in damp habitats Burnet Companion Euclidia glyphica Uncommon/under-recorded. Most frequently recorded from Rutland & east. Burnished Brass Diachrysia chrysitis Common Butterbur Hydraecia petasitis Extinct? VCH & a couple of other records. Under-recorded? Buttoned Snout Hypena rostralis Very rare. VCH record (Buddon Wood) & Ketton Quarry (2015) only Cabbage Moth Mamestra brassicae Fairly common, especially in urban areas.		
Burnet Companion Euclidia glyphica Uncommon/under-recorded. Most frequently recorded from Rutland & east.  Burnished Brass Diachrysia chrysitis Common  Butterbur Hydraecia petasitis Extinct? VCH & a couple of other records. Under-recorded?  Buttoned Snout Hypena rostralis Very rare. VCH record (Buddon Wood) & Ketton Quarry (2015) only  Cabbage Moth Mamestra brassicae Fairly common, especially in urban areas.		
Burnished Brass Diachrysia chrysitis  Butterbur Hydraecia petasitis  Extinct? VCH & a couple of other records. Under-recorded?  Buttoned Snout Hypena rostralis  Very rare. VCH record (Buddon Wood) & Ketton Quarry (2015) only  Cabbage Moth Mamestra brassicae  Fairly common, especially in urban areas.		Uncommon/under-recorded Most frequently recorded from Butland & east
Butterbur Hydraecia petasitis Extinct? VCH & a couple of other records. Under-recorded?  Buttoned Snout Hypena rostralis Very rare. VCH record (Buddon Wood) & Ketton Quarry (2015) only  Cabbage Moth Mamestra brassicae Fairly common, especially in urban areas.		
Buttoned Snout Hypena rostralis Very rare. VCH record (Buddon Wood) & Ketton Quarry (2015) only Cabbage Moth Mamestra brassicae Fairly common, especially in urban areas.		
Cabbage Moth Mamestra brassicae Fairly common, especially in urban areas.		
Campanula rug ⊑upithecia denotata   Kare/localised/under-recorded? Only records: Loddington RTT. 1999 X13 (!)	Campanula Pug Eupithecia denotata	Rare/localised/under-recorded? Only records: Loddington RIT, 1999 x13 (!)

1	15
Campion Sideridis rivularis	Fairly common in gardens
Canary-shouldered Thorn Ennomos alniaria	Common
Centre-barred Sallow Atethmia centrago	Common  Recorded from Dislayarth Ct Wood in 1991 and Clinchem Oversy 2001 02
Chalk Carpet Scotopteryx bipunctaria	Recorded from Pickworth Gt Wood in 1981 and Clipsham Quarry 2001-02.
Chamomile Shark Cucullia chamomillae Chestnut Conistra vaccinii	Uncommon. Recorded only from southern half of County pre 2004.  Common
Chevron Eulithis testata	Declined and now rare/extinct. Only two records since1995.
Chimney Sweeper Odezia atrata	Locally common, particularly in old meadows
Chinese Character Cilix glaucata	Common
Chocolate-tip Clostera curtula	Fairly common. Expanded range & increased in numbers in recent years.
Cinnabar Tyria jacobaeae	Common
Clay Mythimna ferrago	Common
Clay Triple-lines Cyclophora linearia	Rare but increasing in numbers and range, though still localised.
Clifden Nonpareil Catocala fraxini	Very rare migrant. Only approx. 3 very old records.
Cloaked Minor Mesoligia furuncula	Fairly common
Cloaked Pug Eupithecia abietaria	Only 2 records: a Melton garden in 2012 & a Littlethorpe garden in 2015
Clouded Border Lomaspilis marginata	Common
Clouded Brindle Apamea epomidion	Fairly common but local
Clouded Drab Orthosia incerta	Common
Clouded Magpie Abraxas sylvata	Rare. Less than 15 post-2000 records.
Clouded Silver Lomographa temerata	Common.
Clouded-bordered Brindle Apamea crenata	Fairly common
Common Carpet Epirrhoe alternata	Common
Common Emerald Hemithea aestivaria	Common
Common Fan-foot Pechipogo strigilata	Very rare/extinct? Last recorded in 1991. Former status uncertain.
Common Footman Eilema Iurideola	Common
Common Heath Ematurga atomaria	Rare. Only colonies: Ketton Quarry and two other quarries in Rutland.
Common Lutestring Ochropacha duplaris	Locally common in woodland.
Common Marbled Carpet Dysstroma truncata	Common
Common Pug Eupithecia vulgata	Common
Common Quaker Orthosia cerasi	Common
Common Rustic Mesapamea secalis	Common
Common Swift Korscheltellus lupulina	Common
Common Wainscot Mythimna pallens	Common
Common Wave Cabera exanthemata	Common.
Common White Wave Cabera pusaria	Common.
Concolorous Photedes extrema	Rare. Mainly recorded from a handful of sites in Rutland
Confused Apamea furva	Very rare/extinct. Only a handful of records, none since 1994.
Convolvulus Hawk-moth Agrius convolvuli	Generally an uncommon migrant
Copper Underwing Amphipyra pyramidea	Common.
Coronet Craniophora ligustri	Common. Major increase since 2004.
Coxcomb Prominent Ptilodon capucina	Common
Cream Wave Scopula floslactata Cream-bordered Green Pea Earias clorana	Uncommon, particularly in gardens.  Rare & mainly recorded from Rutland & just into Leics.
Crescent Helotropha leucostigma	Rare & localised. Mainly recorded from centre of County (Soar valley?).
Crimson Speckled Utetheisa pulchella	Very rare migrant.
Currant Clearwing Synanthedon tipuliformis	Uncommon/under-recorded (can be common at pheromone lures)
Currant Pug Eupithecia assimilata	Common, but declining?
Dark Arches Apamea monoglypha	Common.
Dark Brocade Mniotype adusta	Status uncertain. Only a handful of records in 1990's, few confirmed.
Dark Chestnut Conistra ligula	Fairly common
Dark Dagger Acronicta tridens	Uncommon/under-recorded (gen det needed)? Fewer records since 1990.
Dark Marbled Carpet Dysstroma citrata	Fairly common
Dark Spectacle Abrostola triplasia	Rare. Only a handful of confirmed records in recent years. (
Dark Spinach Pelurga comitata	Locally common, particularly in Charnwood, but declining?
Dark Sword-grass Agrotis ipsilon	Fairly common migrant
Dark Umber Philereme transversata	Fairly common and more widespread than T. dubitata & P. vetulata.
Dark-barred Twin-spot Carpet Xanthorhoe ferrugata	Uncommon and apparently declining in recent years
Death's-head Hawk-moth Acherontia atropos	Uncommon migrant
December Moth Poecilocampa populi	Common
Deep-brown Dart Aporophyla lutulenta	Uncommon.
Delicate Mythimna vitellina	Rare migrant. Only 10 recent records
Devon Carpet Lampropteryx otregiata	Very rare, only one record: Loddington RIT (2004)
Dingy Footman Eilema griseola	Fairly common.
Dingy Shears Apterogenum ypsillon	Fairly common.
Dingy Shell Euchoeca nebulata	Uncommon and local in damp habitats
Dog's Tooth Lacanobia suasa	Very rare & localised. Post 1982, mainly recorded from Loughbo' Meadows
Dot Moth Melanchra persicariae	Common
Dotted Border Agriopis marginaria	Common
Dotted Chestnut Conistra rubiginea	Rare. VCH + handful of records since 2012. Expanding range?
Dotted Clay Xestia baja	Locally common in Charnwood Forest. Rare elsewhere. Declining?
Dotted Fan-foot Macrochilo cribrumalis	Single record from Rutland Water Rothamsted Trap in 2011
D (1 1 D (1 D) 1 1 1 1	Variable. Fairly common in 1980's, now very rare.
Dotted Rustic Rhyacia simulans	
Double Dart Graphiphora augur	Until recently, fairly common but now seems to be declining.
Double Dart Graphiphora augur Double Lobed Apamea ophiogramma	Fairly common in damp habitats. Few records from extreme west of County.
Double Dart Graphiphora augur Double Lobed Apamea ophiogramma Double Square-spot Xestia triangulum	Fairly common in damp habitats. Few records from extreme west of County.  Common
Double Dart Graphiphora augur Double Lobed Apamea ophiogramma Double Square-spot Xestia triangulum Double-striped Pug Gymnoscelis rufifasciata	Fairly common in damp habitats. Few records from extreme west of County.  Common  Common
Double Dart Graphiphora augur Double Lobed Apamea ophiogramma Double Square-spot Xestia triangulum Double-striped Pug Gymnoscelis rufifasciata Drinker Euthrix potatoria	Fairly common in damp habitats. Few records from extreme west of County.  Common  Common  Common
Double Dart Graphiphora augur Double Lobed Apamea ophiogramma Double Square-spot Xestia triangulum Double-striped Pug Gymnoscelis rufifasciata Drinker Euthrix potatoria Dun-bar Cosmia trapezina	Fairly common in damp habitats. Few records from extreme west of County.  Common  Common  Common  Common.
Double Dart Graphiphora augur Double Lobed Apamea ophiogramma Double Square-spot Xestia triangulum Double-striped Pug Gymnoscelis rufifasciata Drinker Euthrix potatoria	Fairly common in damp habitats. Few records from extreme west of County.  Common  Common  Common

Γ= · - · · · · · · · · · · · · · · · · ·	
Dusky Thorn Ennomos fuscantaria	Fairly common, but declining
Dusky-lemon Sallow Cirrhia gilvago	Uncommon & local. Declined in last 10 years
Dwarf Cream Wave Idaea fuscovenosa  Dwarf Pug Eupithecia tantillaria	Uncommon and but expanding its range in county & increasing in numbers.
	Fairly common.  Uncommon & local. Most frequently recorded from Charnwood Forest.
Ear Moth Amphipoea oculea  Earias vittella	Nationally rare. Only one VC55 record, from a Wigston garden in 2006.
Early Grey Xylocampa areola	Common
Early Moth Theria primaria	Uncommon. Declining and/or under-recorded?
Early Thorn Selenia dentaria	Common
Early Tooth-striped Trichopteryx carpinata	Rare & local in western half of county only, but commoner in last 10 years
Elephant Hawk-moth Deilephila elpenor	Common
Emperor Moth Saturnia pavonia	Rare. Recent records are from the (north) east of the County/Rutland
Engrailed Ectropis crepuscularia	Common.
Eyed Hawk-moth Smerinthus ocellata	Fairly common
False Mocha Cyclophora porata	Very rare, but four records from Rutland in 2004-2009.
Fan-foot Herminia tarsipennalis	Common
Feathered Gothic Tholera decimalis	Fairly common.
Feathered Ranunculus Polymixis lichenea	One record only: Preston 1998. Vagrant/migrant?
Feathered Thorn Colotois pennaria	Common
Fen Wainscot Arenostola phragmitidis	Uncommon mainly recorded from a handful of localities.
Fern Horisme tersata	Rare and localised to a handful of sites with Traveller's Joy
Festoon Apoda limacodes	Very rare. Only recorded from Ketton Quarry in 2008 & 2015
Figure of Eight Diloba caeruleocephala	Uncommon & declining
Figure of Eighty Tethea ocularis	Common
Flame Axylia putris	Common
Flame Carpet Xanthorhoe designata	Fairly common in woodland and regular in many gardens
Flame Shoulder Ochropleura plecta	Common
Flounced Chestnut Agrochola helvola	Uncommon. Rarely recorded outside of Charnwood Forest.
Flounced Rustic Luperina testacea	Common
Forester Adscita statices	Very local & uncommon
Four-dotted Footman Cybosia mesomella	Uncommon. Rarely recorded outside of Rutland
Four-spotted Footman Lithosia quadra	Very rare migrant.
Four-spotted Tyta luctuosa	Very Rare. A colony in Ketton area, plus a few other records.
Fox Moth Macrothylacia rubi	Very rare. Only know from Charnwood Lodge and Humberstone (2015)
Foxglove Pug Eupithecia pulchellata	Common
Freyer's Pug Eupithecia intricata	Common. Increased dramatically in last 20 years.
Frosted Green Polyploca ridens	Rare. Most recent records are from Rutland woodland sites.
Frosted Orange Gortyna flavago	Fairly common. Declining?
Galium Carpet Epirrhoe galiata	Uncommon and localised
Garden Carpet Xanthorhoe fluctuata	Common
Garden Dart Euxoa nigricans	Rare & declining
Garden Tiger Arctia caja	Formerly common, but declined dramatically & now uncommon/rare.
Gem Nycterosea obstipata	Uncommon migrant
Ghost Moth Hepialus humuli Glaucous Shears Papestra biren	Fairly common  Very rare & restricted to Charnwood Forest. Few recent records.
Goat Moth Cossus cossus Gold Spangle Autographa bractea	Now very rare. Only one recent record: Brown's Hill Quarry, 1990  Very rare. Only a handful of records in last 10 years. Declining?
Gold Spot Plusia festucae	Uncommon & local. Found mainly in the west.
Gold Swift Phymatopus hecta	Uncommon, especially outside of woodland
Golden Plusia Polychrysia moneta	Fairly common in gardens in 1970's/1980's, less so now.
Golden Twin-spot Chrysodeixis chalcites	Very rare migrant. Only one record from a Ravenstone garden in 2006.
Golden-rod Brindle Xylena solidaginis	Probably extinct. Last record: Charnwood Lodge, 1980.
Golden-rod Pug Eupithecia virgaureata	Uncommon. Mainly recorded from western half of county.
Gothic Naenia typica	Common
Grass Emerald Pseudoterpna pruinata	Rare. Only recorded from Sapcote and a single site in Rutland since 2000.
Grass Rivulet Perizoma albulata	Rare & localised. Mainly recorded from Charnwood Forest
Grass Wave Perconia strigillaria	Very rare. Ketton Quarry the only known colony
Great Brocade Eurois occulta	Rare migrant.
Green Arches Anaplectoides prasina	Locally common in wooded areas.
Green Carpet Colostygia pectinataria	Common
	Common
Green Pug Pasiphila rectangulata	Common
Green Pug Pasiphila rectangulata Green Silver-lines Pseudoips prasinana	Fairly common, particularly in west of County.
Green Silver-lines Pseudoips prasinana Green-brindled Crescent Allophyes oxyacanthae	Fairly common, particularly in west of County.  Common
Green Silver-lines Pseudoips prasinana Green-brindled Crescent Allophyes oxyacanthae Grey Arches Polia nebulosa	Fairly common, particularly in west of County.  Common  Uncommon and local, mainly recorded from Charnwood Forest
Green Silver-lines Pseudoips prasinana Green-brindled Crescent Allophyes oxyacanthae Grey Arches Polia nebulosa Grey Birch Aethalura punctulata	Fairly common, particularly in west of County.  Common  Uncommon and local, mainly recorded from Charnwood Forest  Uncommon. A woodland species commonest in west of County.
Green Silver-lines Pseudoips prasinana Green-brindled Crescent Allophyes oxyacanthae Grey Arches Polia nebulosa Grey Birch Aethalura punctulata Grey Chi Antitype chi	Fairly common, particularly in west of County.  Common  Uncommon and local, mainly recorded from Charnwood Forest  Uncommon. A woodland species commonest in west of County.  Rare. Declining?
Green Silver-lines Pseudoips prasinana Green-brindled Crescent Allophyes oxyacanthae Grey Arches Polia nebulosa Grey Birch Aethalura punctulata Grey Chi Antitype chi Grey Dagger Acronicta psi	Fairly common, particularly in west of County.  Common  Uncommon and local, mainly recorded from Charnwood Forest  Uncommon. A woodland species commonest in west of County.  Rare. Declining?  Common.
Green Silver-lines Pseudoips prasinana Green-brindled Crescent Allophyes oxyacanthae Grey Arches Polia nebulosa Grey Birch Aethalura punctulata Grey Chi Antitype chi Grey Dagger Acronicta psi Grey Pine Carpet Thera obeliscata	Fairly common, particularly in west of County.  Common  Uncommon and local, mainly recorded from Charnwood Forest  Uncommon. A woodland species commonest in west of County.  Rare. Declining?  Common.  Common
Green Silver-lines Pseudoips prasinana Green-brindled Crescent Allophyes oxyacanthae Grey Arches Polia nebulosa Grey Birch Aethalura punctulata Grey Chi Antitype chi Grey Dagger Acronicta psi Grey Pine Carpet Thera obeliscata Grey Pug Eupithecia subfuscata	Fairly common, particularly in west of County.  Common  Uncommon and local, mainly recorded from Charnwood Forest  Uncommon. A woodland species commonest in west of County.  Rare. Declining?  Common.  Common  Common
Green Silver-lines Pseudoips prasinana Green-brindled Crescent Allophyes oxyacanthae Grey Arches Polia nebulosa Grey Birch Aethalura punctulata Grey Chi Antitype chi Grey Dagger Acronicta psi Grey Pine Carpet Thera obeliscata Grey Pug Eupithecia subfuscata Grey Shoulder-knot Lithophane ornitopus	Fairly common, particularly in west of County.  Common  Uncommon and local, mainly recorded from Charnwood Forest  Uncommon. A woodland species commonest in west of County.  Rare. Declining?  Common.  Common  Common  Fairly common.
Green Silver-lines Pseudoips prasinana Green-brindled Crescent Allophyes oxyacanthae Grey Arches Polia nebulosa Grey Birch Aethalura punctulata Grey Chi Antitype chi Grey Dagger Acronicta psi Grey Pine Carpet Thera obeliscata Grey Pug Eupithecia subfuscata Grey Shoulder-knot Lithophane ornitopus Gypsy Moth Lymantria dispar	Fairly common, particularly in west of County.  Common  Uncommon and local, mainly recorded from Charnwood Forest  Uncommon. A woodland species commonest in west of County.  Rare. Declining?  Common.  Common  Common  Fairly common.  Extinct
Green Silver-lines Pseudoips prasinana Green-brindled Crescent Allophyes oxyacanthae Grey Arches Polia nebulosa Grey Birch Aethalura punctulata Grey Chi Antitype chi Grey Dagger Acronicta psi Grey Pine Carpet Thera obeliscata Grey Pug Eupithecia subfuscata Grey Shoulder-knot Lithophane ornitopus Gypsy Moth Lymantria dispar Haworth's Pug Eupithecia haworthiata	Fairly common, particularly in west of County.  Common  Uncommon and local, mainly recorded from Charnwood Forest  Uncommon. A woodland species commonest in west of County.  Rare. Declining?  Common.  Common  Common  Fairly common.  Extinct  Rare and localised. Recorded from a handful of sites only.
Green Silver-lines Pseudoips prasinana Green-brindled Crescent Allophyes oxyacanthae Grey Arches Polia nebulosa Grey Birch Aethalura punctulata Grey Chi Antitype chi Grey Dagger Acronicta psi Grey Pine Carpet Thera obeliscata Grey Pug Eupithecia subfuscata Grey Shoulder-knot Lithophane ornitopus Gypsy Moth Lymantria dispar Haworth's Pug Eupithecia haworthiata Heart & Club Agrotis clavis	Fairly common, particularly in west of County.  Common  Uncommon and local, mainly recorded from Charnwood Forest  Uncommon. A woodland species commonest in west of County.  Rare. Declining?  Common.  Common  Common  Fairly common.  Extinct  Rare and localised. Recorded from a handful of sites only.  Uncommon but increasing at present time.
Green Silver-lines Pseudoips prasinana Green-brindled Crescent Allophyes oxyacanthae Grey Arches Polia nebulosa Grey Birch Aethalura punctulata Grey Chi Antitype chi Grey Dagger Acronicta psi Grey Pine Carpet Thera obeliscata Grey Pug Eupithecia subfuscata Grey Shoulder-knot Lithophane ornitopus Gypsy Moth Lymantria dispar Haworth's Pug Eupithecia haworthiata Heart & Club Agrotis clavis Heart & Dart Agrotis exclamationis	Fairly common, particularly in west of County.  Common  Uncommon and local, mainly recorded from Charnwood Forest  Uncommon. A woodland species commonest in west of County.  Rare. Declining?  Common.  Common  Common  Fairly common.  Extinct  Rare and localised. Recorded from a handful of sites only.  Uncommon but increasing at present time.  Common, especially in gardens
Green Silver-lines Pseudoips prasinana Green-brindled Crescent Allophyes oxyacanthae Grey Arches Polia nebulosa Grey Birch Aethalura punctulata Grey Chi Antitype chi Grey Dagger Acronicta psi Grey Pine Carpet Thera obeliscata Grey Pug Eupithecia subfuscata Grey Shoulder-knot Lithophane ornitopus Gypsy Moth Lymantria dispar Haworth's Pug Eupithecia haworthiata Heart & Club Agrotis clavis Heart & Dart Agrotis exclamationis Heath Rustic Xestia agathina	Fairly common, particularly in west of County.  Common  Uncommon and local, mainly recorded from Charnwood Forest  Uncommon. A woodland species commonest in west of County.  Rare. Declining?  Common.  Common  Common  Fairly common.  Extinct  Rare and localised. Recorded from a handful of sites only.  Uncommon but increasing at present time.  Common, especially in gardens  Rare. Only two known colonies: Warren Hills & Newfield Colliery.
Green Silver-lines Pseudoips prasinana Green-brindled Crescent Allophyes oxyacanthae Grey Arches Polia nebulosa Grey Birch Aethalura punctulata Grey Chi Antitype chi Grey Dagger Acronicta psi Grey Pine Carpet Thera obeliscata Grey Pug Eupithecia subfuscata Grey Shoulder-knot Lithophane ornitopus Gypsy Moth Lymantria dispar Haworth's Pug Eupithecia haworthiata Heart & Club Agrotis clavis Heart & Dart Agrotis exclamationis Heath Rustic Xestia agathina Hebrew Character Orthosia gothica	Fairly common, particularly in west of County.  Common  Uncommon and local, mainly recorded from Charnwood Forest  Uncommon. A woodland species commonest in west of County.  Rare. Declining?  Common.  Common  Common  Fairly common.  Extinct  Rare and localised. Recorded from a handful of sites only.  Uncommon but increasing at present time.  Common, especially in gardens  Rare. Only two known colonies: Warren Hills & Newfield Colliery.  Common
Green Silver-lines Pseudoips prasinana Green-brindled Crescent Allophyes oxyacanthae Grey Arches Polia nebulosa Grey Birch Aethalura punctulata Grey Chi Antitype chi Grey Dagger Acronicta psi Grey Pine Carpet Thera obeliscata Grey Pug Eupithecia subfuscata Grey Shoulder-knot Lithophane ornitopus Gypsy Moth Lymantria dispar Haworth's Pug Eupithecia haworthiata Heart & Club Agrotis clavis Heart & Dart Agrotis exclamationis Heath Rustic Xestia agathina Hebrew Character Orthosia gothica Hedge Rustic Tholera cespitis	Fairly common, particularly in west of County.  Common  Uncommon and local, mainly recorded from Charnwood Forest  Uncommon. A woodland species commonest in west of County.  Rare. Declining?  Common.  Common  Common  Fairly common.  Extinct  Rare and localised. Recorded from a handful of sites only.  Uncommon but increasing at present time.  Common, especially in gardens  Rare. Only two known colonies: Warren Hills & Newfield Colliery.  Common  Uncommon and local. Mainly recorded in Charnwood Forest.
Green Silver-lines Pseudoips prasinana Green-brindled Crescent Allophyes oxyacanthae Grey Arches Polia nebulosa Grey Birch Aethalura punctulata Grey Chi Antitype chi Grey Dagger Acronicta psi Grey Pine Carpet Thera obeliscata Grey Pug Eupithecia subfuscata Grey Shoulder-knot Lithophane ornitopus Gypsy Moth Lymantria dispar Haworth's Pug Eupithecia haworthiata Heart & Club Agrotis clavis Heart & Dart Agrotis exclamationis Heath Rustic Xestia agathina Hebrew Character Orthosia gothica Hedge Rustic Tholera cespitis Herald Scoliopteryx libatrix	Fairly common, particularly in west of County.  Common  Uncommon and local, mainly recorded from Charnwood Forest  Uncommon. A woodland species commonest in west of County.  Rare. Declining?  Common.  Common  Common  Fairly common.  Extinct  Rare and localised. Recorded from a handful of sites only.  Uncommon but increasing at present time.  Common, especially in gardens  Rare. Only two known colonies: Warren Hills & Newfield Colliery.  Common  Uncommon and local. Mainly recorded in Charnwood Forest.  Common
Green Silver-lines Pseudoips prasinana Green-brindled Crescent Allophyes oxyacanthae Grey Arches Polia nebulosa Grey Birch Aethalura punctulata Grey Chi Antitype chi Grey Dagger Acronicta psi Grey Pine Carpet Thera obeliscata Grey Pug Eupithecia subfuscata Grey Shoulder-knot Lithophane ornitopus Gypsy Moth Lymantria dispar Haworth's Pug Eupithecia haworthiata Heart & Club Agrotis clavis Heart & Dart Agrotis exclamationis Heath Rustic Xestia agathina Hebrew Character Orthosia gothica Hedge Rustic Tholera cespitis Herald Scoliopteryx libatrix Hoary Footman Eilema caniola	Fairly common, particularly in west of County.  Common  Uncommon and local, mainly recorded from Charnwood Forest  Uncommon. A woodland species commonest in west of County.  Rare. Declining?  Common.  Common  Common  Fairly common.  Extinct  Rare and localised. Recorded from a handful of sites only.  Uncommon but increasing at present time.  Common, especially in gardens  Rare. Only two known colonies: Warren Hills & Newfield Colliery.  Common  Uncommon and local. Mainly recorded in Charnwood Forest.  Common  Rare. Recorded from 4 gardens and may be expanding (
Green Silver-lines Pseudoips prasinana Green-brindled Crescent Allophyes oxyacanthae Grey Arches Polia nebulosa Grey Birch Aethalura punctulata Grey Chi Antitype chi Grey Dagger Acronicta psi Grey Pine Carpet Thera obeliscata Grey Pug Eupithecia subfuscata Grey Shoulder-knot Lithophane ornitopus Gypsy Moth Lymantria dispar Haworth's Pug Eupithecia haworthiata Heart & Club Agrotis clavis Heart & Dart Agrotis exclamationis Heath Rustic Xestia agathina Hebrew Character Orthosia gothica Hedge Rustic Tholera cespitis Herald Scoliopteryx libatrix	Fairly common, particularly in west of County.  Common  Uncommon and local, mainly recorded from Charnwood Forest  Uncommon. A woodland species commonest in west of County.  Rare. Declining?  Common.  Common  Common  Fairly common.  Extinct  Rare and localised. Recorded from a handful of sites only.  Uncommon but increasing at present time.  Common, especially in gardens  Rare. Only two known colonies: Warren Hills & Newfield Colliery.  Common  Uncommon and local. Mainly recorded in Charnwood Forest.  Common

Iron Prominent Notodonta dromedarius  July Belle Scotopteryx luridata  July Highflier Hydriomena furcata  Co Juniper Carpet Thera juniperata  No Juniper Pug Eupithecia pusillata  Knot Grass Acronicta rumicis  Lackey Malacosoma neustria  Lappet Gastropacha quercifolia  Ve Larch Pug Eupithecia lariciata  Un Large Emerald Geometra papilionaria  Fa Large Nutmeg Apamea anceps  Co Large Ranunculus Polymixis flavicincta  Large Red-belted Clearwing Synanthedon culiciformis  Ve Large Thorn Ennomos autumnaria  Ra Large Twin-spot Carpet Xanthorhoe quadrifasiata	common care/extinct? Not recorded since 1987. common com fairly common in gardens, though not as common in 1980s-90s. airly common, especially in gardens. cocally common, particularly in west of County. airly common cery rare. common/under-recorded common in Rutland & the east, rarer in west of county. Increasing? cocally common in Rutland (and just into Leics.). Unrecorded elsewhere. cery rare/under-recorded. Last record: Spring Wood, 1984. care. Only 5 post-VCH records: Barrowden 2000-08 & Loddington RIT 2010 cow fairly common. common and local. Most records from eastern half of county common common common common common cery rare/extinct. Possible confusion with S. luridata; checking required.
July Belle Scotopteryx Iuridata  July Highflier Hydriomena furcata  Co Juniper Carpet Thera juniperata  No Juniper Pug Eupithecia pusillata  Knot Grass Acronicta rumicis  Lackey Malacosoma neustria  Lappet Gastropacha quercifolia  Ve Larch Pug Eupithecia Iariciata  Un Large Emerald Geometra papilionaria  Fa Large Nutmeg Apamea anceps  Co Large Ranunculus Polymixis flavicincta  Large Red-belted Clearwing Synanthedon culiciformis  Ve Large Thorn Ennomos autumnaria  Ra Large Twin-spot Carpet Xanthorhoe quadrifasiata	common low fairly common in gardens, though not as common in 1980s-90s. airly common, especially in gardens. locally common, particularly in west of County. airly common low fairly common, particularly in west of County. airly common locally common in woodland. locally common in Rutland & the east, rarer in west of county. Increasing? locally common in Rutland (and just into Leics.). Unrecorded elsewhere. lery rare/under-recorded. Last record: Spring Wood, 1984. lare. Only 5 post-VCH records: Barrowden 2000-08 & Loddington RIT 2010 low fairly common. locommon and local. Most records from eastern half of county locommon
July Highflier Hydriomena furcata  Juniper Carpet Thera juniperata  Not Juniper Pug Eupithecia pusillata  Knot Grass Acronicta rumicis  Lackey Malacosoma neustria  Lappet Gastropacha quercifolia  Velarch Pug Eupithecia lariciata  Large Emerald Geometra papilionaria  Falarge Nutmeg Apamea anceps  Collarge Ranunculus Polymixis flavicincta  Large Red-belted Clearwing Synanthedon culiciformis  Velarge Thorn Ennomos autumnaria  Ratarge Twin-spot Carpet Xanthorhoe quadrifasiata	common low fairly common in gardens, though not as common in 1980s-90s. lairly common, especially in gardens. locally common, particularly in west of County. lairly common lery rare. locommon/under-recorded lairly common in woodland. locommon in Rutland & the east, rarer in west of county. Increasing? locally common in Rutland (and just into Leics.). Unrecorded elsewhere. lery rare/under-recorded. Last record: Spring Wood, 1984. lare. Only 5 post-VCH records: Barrowden 2000-08 & Loddington RIT 2010 low fairly common. locommon and local. Most records from eastern half of county locommon
Juniper Carpet Thera juniperata Juniper Pug Eupithecia pusillata Knot Grass Acronicta rumicis Lackey Malacosoma neustria Fa Lappet Gastropacha quercifolia Ve Larch Pug Eupithecia lariciata Un Large Emerald Geometra papilionaria Fa Large Nutmeg Apamea anceps Cc Large Ranunculus Polymixis flavicincta Large Red-belted Clearwing Synanthedon culiciformis Ve Large Thorn Ennomos autumnaria Ra Large Twin-spot Carpet Xanthorhoe quadrifasiata	ow fairly common in gardens, though not as common in 1980s-90s. airly common, especially in gardens. ocally common, particularly in west of County. airly common ery rare. incommon/under-recorded airly common in woodland. common in Rutland & the east, rarer in west of county. Increasing? ocally common in Rutland (and just into Leics.). Unrecorded elsewhere. ery rare/under-recorded. Last record: Spring Wood, 1984. tare. Only 5 post-VCH records: Barrowden 2000-08 & Loddington RIT 2010 low fairly common. Incommon and local. Most records from eastern half of county
Juniper Pug Eupithecia pusillata Fa Knot Grass Acronicta rumicis Lo Lackey Malacosoma neustria Fa Lappet Gastropacha quercifolia Ve Larch Pug Eupithecia lariciata Un Large Emerald Geometra papilionaria Fa Large Nutmeg Apamea anceps Cc Large Ranunculus Polymixis flavicincta Lo Large Red-belted Clearwing Synanthedon culiciformis Ve Large Thorn Ennomos autumnaria Ra Large Twin-spot Carpet Xanthorhoe quadrifasiata	airly common, especially in gardens. ocally common, particularly in west of County. airly common ery rare. Incommon/under-recorded airly common in woodland. common in Rutland & the east, rarer in west of county. Increasing? ocally common in Rutland (and just into Leics.). Unrecorded elsewhere. ery rare/under-recorded. Last record: Spring Wood, 1984. lare. Only 5 post-VCH records: Barrowden 2000-08 & Loddington RIT 2010 low fairly common. Incommon and local. Most records from eastern half of county
Knot Grass Acronicta rumicis Lackey Malacosoma neustria Fa Lappet Gastropacha quercifolia Ve Larch Pug Eupithecia lariciata Un Large Emerald Geometra papilionaria Fa Large Nutmeg Apamea anceps Cc Large Ranunculus Polymixis flavicincta Large Red-belted Clearwing Synanthedon culiciformis Ve Large Thorn Ennomos autumnaria Ra Large Twin-spot Carpet Xanthorhoe quadrifasiata	ocally common, particularly in west of County.  airly common  ery rare. Incommon/under-recorded  airly common in woodland.  common in Rutland & the east, rarer in west of county. Increasing?  ocally common in Rutland (and just into Leics.). Unrecorded elsewhere.  ery rare/under-recorded. Last record: Spring Wood, 1984.  lare. Only 5 post-VCH records: Barrowden 2000-08 & Loddington RIT 2010  low fairly common.  Incommon and local. Most records from eastern half of county  common
Lackey Malacosoma neustria  Lappet Gastropacha quercifolia  Ve  Larch Pug Eupithecia Iariciata  Un  Large Emerald Geometra papilionaria  Fa  Large Nutmeg Apamea anceps  Cc  Large Ranunculus Polymixis flavicincta  Large Red-belted Clearwing Synanthedon culiciformis  Ve  Large Thorn Ennomos autumnaria  Ra  Large Twin-spot Carpet Xanthorhoe quadrifasiata	airly common ery rare. Incommon/under-recorded airly common in woodland. Incommon in Rutland & the east, rarer in west of county. Increasing? Incommon in Rutland (and just into Leics.). Unrecorded elsewhere. Incommon in Rutland (and just into Leics.). Unrecorded elsewhere. Incommon in Rutland (and just into Leics.). Unrecorded elsewhere. Incommon in Rutland (and just into Leics.). Unrecorded elsewhere. Incommon in Rutland (and just into Leics.). Unrecorded elsewhere. Incommon in Rutland (and just into Leics.). Unrecorded elsewhere. Incommon in Rutland (and just into Leics.). Unrecorded elsewhere. Incommon in Rutland (and just into Leics.). Unrecorded elsewhere. Incommon in Rutland (and just into Leics.). Unrecorded elsewhere. Incommon in Rutland (and just into Leics.). Unrecorded elsewhere. Incommon in Rutland (and just into Leics.). Unrecorded elsewhere. Incommon in Rutland (and just into Leics.). Unrecorded elsewhere. Incommon in Rutland (and just into Leics.). Unrecorded elsewhere. Incommon in Rutland (and just into Leics.). Unrecorded elsewhere. Incommon in Rutland (and just into Leics.). Unrecorded elsewhere. Incommon in Rutland (and just into Leics.). Unrecorded elsewhere. Incommon in Rutland (and just into Leics.). Unrecorded elsewhere. Incommon in Rutland (and just into Leics.). Unrecorded elsewhere. Incommon in Rutland (and just into Leics.). Unrecorded elsewhere. Incommon in Rutland (and just into Leics.). Unrecorded elsewhere. Incommon in Rutland (and just into Leics.). Unrecorded elsewhere. Incommon in Rutland (and just into Leics.). Unrecorded elsewhere. Incommon in Rutland (and just into Leics.). Unrecorded elsewhere. Incommon in Rutland (and just into Leics.). Unrecorded elsewhere. Incommon in Rutland (and just into Leics.). Unrecorded elsewhere. Incommon in Rutland (and just into Leics.). Unrecorded elsewhere. Incommon in Rutland (and just into Leics.). Unrecorded elsewhere. Incommon in Rutland (and just into Leics.). Unrecorded elsewhere. Incommon in Rutland (and just into Leics.). Unrecorded e
Lappet Gastropacha quercifolia  Larch Pug Eupithecia Iariciata  Un  Large Emerald Geometra papilionaria  Fa  Large Nutmeg Apamea anceps  Co  Large Ranunculus Polymixis flavicincta  Large Red-belted Clearwing Synanthedon culiciformis  Ve  Large Thorn Ennomos autumnaria  Ratarge Twin-spot Carpet Xanthorhoe quadrifasiata	ery rare. Incommon/under-recorded airly common in woodland. Incommon in Rutland & the east, rarer in west of county. Increasing? Incommon in Rutland (and just into Leics.). Unrecorded elsewhere. Incommon in Rutland (and just into Leics.). Unrecorded elsewhere. Incommon in Rutland (and just into Leics.). Unrecorded elsewhere. Incommon in Rutland (and just into Leics.). Unrecorded elsewhere. Incommon in Rutland (and just into Leics.). Unrecorded elsewhere. Incommon in Rutland (and just into Leics.). Unrecorded elsewhere. Incommon in Rutland (and just into Leics.). Unrecorded elsewhere. Incommon in Rutland (and just into Leics.). Unrecorded elsewhere. Incommon in Rutland (and just into Leics.). Unrecorded elsewhere. Incommon in Rutland (and just into Leics.). Unrecorded elsewhere. Incommon in Rutland (and just into Leics.). Unrecorded elsewhere. Incommon in Rutland (and just into Leics.). Unrecorded elsewhere. Incommon in Rutland (and just into Leics.). Unrecorded elsewhere. Incommon in Rutland (and just into Leics.). Unrecorded elsewhere. Incommon in Rutland (and just into Leics.). Unrecorded elsewhere. Incommon in Rutland (and just into Leics.). Unrecorded elsewhere. Incommon in Rutland (and just into Leics.). Unrecorded elsewhere. Incommon in Rutland (and just into Leics.). Unrecorded elsewhere. Incommon in Rutland (and just into Leics.). Unrecorded elsewhere. Incommon in Rutland (and just into Leics.). Unrecorded elsewhere. Incommon in Rutland (and just into Leics.). Unrecorded elsewhere. Incommon in Rutland (and just into Leics.). Unrecorded elsewhere. Incommon in Rutland (and just into Leics.). Unrecorded elsewhere. Incommon in Rutland (and just into Leics.). Unrecorded elsewhere. Incommon in Rutland (and just into Leics.). Unrecorded elsewhere. Incommon in Rutland (and just into Leics.). Unrecorded elsewhere. Incommon in Rutland (and just into Leics.). Unrecorded elsewhere. Incommon in Rutland (and just into Leics.). Unrecorded elsewhere. Incommon in Rutland (and just into Leics.). Unrecorded elsewhere. Inc
Larch Pug Eupithecia Iariciata Un Large Emerald Geometra papilionaria Fa Large Nutmeg Apamea anceps Cc Large Ranunculus Polymixis flavicincta Lo Large Red-belted Clearwing Synanthedon culiciformis Ve Large Thorn Ennomos autumnaria Ra Large Twin-spot Carpet Xanthorhoe quadrifasiata No	ncommon/under-recorded airly common in woodland. common in Rutland & the east, rarer in west of county. Increasing? cocally common in Rutland (and just into Leics.). Unrecorded elsewhere. ery rare/under-recorded. Last record: Spring Wood, 1984. care. Only 5 post-VCH records: Barrowden 2000-08 & Loddington RIT 2010 low fairly common. Incommon and local. Most records from eastern half of county common
Large Emerald Geometra papilionaria Fa Large Nutmeg Apamea anceps Co Large Ranunculus Polymixis flavicincta Lo Large Red-belted Clearwing Synanthedon culiciformis Ve Large Thorn Ennomos autumnaria Ra Large Twin-spot Carpet Xanthorhoe quadrifasiata No	airly common in woodland. common in Rutland & the east, rarer in west of county. Increasing? cocally common in Rutland (and just into Leics.). Unrecorded elsewhere. ery rare/under-recorded. Last record: Spring Wood, 1984. care. Only 5 post-VCH records: Barrowden 2000-08 & Loddington RIT 2010 low fairly common. Incommon and local. Most records from eastern half of county common
Large Nutmeg Apamea anceps Co Large Ranunculus Polymixis flavicincta Lo Large Red-belted Clearwing Synanthedon culiciformis Ve Large Thorn Ennomos autumnaria Ra Large Twin-spot Carpet Xanthorhoe quadrifasiata No	common in Rutland & the east, rarer in west of county. Increasing? cocally common in Rutland (and just into Leics.). Unrecorded elsewhere. ery rare/under-recorded. Last record: Spring Wood, 1984. care. Only 5 post-VCH records: Barrowden 2000-08 & Loddington RIT 2010 low fairly common. Incommon and local. Most records from eastern half of county common
Large Ranunculus Polymixis flavicincta Lo Large Red-belted Clearwing Synanthedon culiciformis Ve Large Thorn Ennomos autumnaria Ra Large Twin-spot Carpet Xanthorhoe quadrifasiata No	ocally common in Rutland (and just into Leics.). Unrecorded elsewhere. ery rare/under-recorded. Last record: Spring Wood, 1984. lare. Only 5 post-VCH records: Barrowden 2000-08 & Loddington RIT 2010 low fairly common. Incommon and local. Most records from eastern half of county common
Large Red-belted Clearwing Synanthedon culiciformis  Ve Large Thorn Ennomos autumnaria Ra Large Twin-spot Carpet Xanthorhoe quadrifasiata No	ery rare/under-recorded. Last record: Spring Wood, 1984. lare. Only 5 post-VCH records: Barrowden 2000-08 & Loddington RIT 2010 low fairly common. Incommon and local. Most records from eastern half of county common
Large Thorn Ennomos autumnaria Ra Large Twin-spot Carpet Xanthorhoe quadrifasiata No	lare. Only 5 post-VCH records: Barrowden 2000-08 & Loddington RIT 2010 ow fairly common. Incommon and local. Most records from eastern half of county common common
Large Twin-spot Carpet Xanthorhoe quadrifasiata No	ow fairly common. Incommon and local. Most records from eastern half of county common common
	ncommon and local. Most records from eastern half of county common common
	common common
	ommon
	ncommon and local, mainly recorded from eastern half of county
	ocally common
	ncommon, but recent records indicate that this species is increasing.
	airly common.
	common
	common
	airly common/under-recorded (few gen-det confirmed ids)
	are and apparently declining. (
Lesser Swallow Prominent Pheosia gnoma Co	ommon, especially in habitats with birch
	are/uncommon. Fewer records recently, mainly from west of county.
	ommon
Lesser-spotted Pinion Cosmia affinis Ra	are. Only recent records are from a handful of sites in Rutland.
Light Arches Apamea lithoxylaea Co	ommon.
Light Brocade Lacanobia w-latinum Lo	ocally common in Rutland, rare elsewhere
Light Emerald Campaea margaritaria Co	ommon.
Light Orange Underwing Boudinotiana notha Ve	ery rare/extinct/unrecorded. Last record: Burbage Wood 1940's?
Lilac Beauty Apeira syringaria Fa	airly common
	airly common
Lime-speck Pug Eupithecia centaureata Co	ommon
	Now considered a sub-species of Wormwood Pug. Charnwood Lodge)
Little Emerald Jodis lactearia Fa	airly common but localised?
	are/under-recorded. Very few recent records.
,	ommon
	eclining and now very rare/extinct. Last record: Markfield, 1990.
	ommon
	ncommon and probably declining
	airly common in gardens
	common
	ncommon, but increasing and now turning-up in gardens.
	common in Rutland gardens, very rare elsewhere lare and restricted to Rutland, with some expansion into east Leics.
	•
1 0 1	Incommon/under-recorded
	ocally common in west of County common, especially in built-up areas.
	ocally common in Charnwood Forest. Very rare elsewhere
	Incommon and local in gardens in Rutland & southern half of County.
	ery Rare. VCH + recent records from Loddington RIT.
	common (but few gen-det confirmed ids)
	ocally common in Rutland, uncommon/rare elsewhere
	common
	are. Few recent records. Dayflying & under-recorded?
	ncommon. Mainly found in damp habitats in west of County
	ncommon and local. Most frequently recorded from Rutland
	airly common
	common
Miller Acronicta leporina Lo	ocally common, particularly in west of County.
	airly common in woodland.
Mother Shipton Euclidia mi Un	ncommon/under-recorded.
	ommon
Mottled Grey Colostygia multistrigaria Lo	ocally common in Charnwood Forest. Unrecorded elsewhere
U I U	ommon
	ommon, especially in gardens.
	ommon
	ommon.
	airly common/under-recorded (most commonly recorded as larvae)
	ncommon.
	airly common. Declining?
	xtinct. VCH only.
	airly common. Under-recorded?
0 0 1	ncommon.
Netted Pug Eupithecia venosata Ra	are. Rutland only.

Ni Moth Trishonlusia ni	Vanurara migrant Only 2 records Laignator (4002) 9 Quaniharayah (2015)
Ni Moth Trichoplusia ni Northern Drab Orthosia opima	Very rare migrant. Only 2 records Leicester (1982) & Queniborough (2015)  Extinct? Last record: Pillings Lock, 1985.
Northern Rustic Standfussiana lucernea	Very rare/extinct? Only record: Charnwood Lodge, 1965.
Northern Spinach Eulithis populata	Uncommon. Mainly recorded from Charnwood, but declining.
Northern Winter Moth Operophtera fagata	Uncommon, mainly recorded from west of county.
November Moth Epirrita dilutata	Common (though few records confirmed by gen det)
Nutmeg Anarta trifolii	Fairly common
Nut-tree Tussock Colocasia coryli	VCH record from Buddon Wood.
Oak Beauty Biston strataria	Fairly common
Oak Eggar Lasiocampa quercus	Uncommon. Mainly recorded from southern half of county.
Oak Hook-tip Watsonalla binaria	Common
Oak Lutestring Cymatophorina diluta	Very rare. Only a handful of recent records but a colony at Spring Wood
Oak Nycteoline Nycteola revayana	Uncommon, but increasing since mid-1990s. Probably under-recorded.
Oak-tree Pug Eupithecia dodoneata Oblique Carpet Orthonama vittata	Fairly common  Very rare. A 2006 record from Ravenstone was the first for 30+ years.
Obscure Wainscot Leucania obsoleta	Very rare. Only known from Rutland Water & Eyebrook Reservoir
Ochreous Pug Eupithecia indigata	Uncommon
Old Lady Mormo maura	Fairly common
Oleander Hawk-moth Daphnis nerii	Very rare migrant.
Olive Ipimorpha subtusa	Locally fairly common. Increasing?
Orange Footman Eilema sororcula	Common. First recorded in 2003, & increased rapidly since then.
Orange Moth Angerona prunaria	Extinct? Only a couple of probably doubtful post VCH records
Orange Sallow Tiliacea citrago	Uncommon & local.
Orange Swift Triodia sylvina	Common
Orange Underwing Archiearis parthenias	Uncommon, but under-recorded
Orange-tailed Clearwing Synanthedon	Rare/under-recorded. Only one record: Tinwell, 27/06/05 (to lure)
andrenaeformis	
Pale Brindled Beauty Phigalia pilosaria	Fairly common
Pale Eggar Trichiura crataegi	Fairly common. Declining.
Pale Mottled Willow Caradrina clavipalpis	Common
Pale November Moth Epirrita christyi Pale Oak Beauty Hypomecis punctinalis	Less common than E. dilutata? (few records confirmed by gen det) Uncommon/rare & mainly recorded from Rutland woodland sites, but increasing
Pale Pinion Lithophane socia	Fairly common. Only arrived in county in 1996
Pale Prominent Pterostoma palpina	Common
Pale Shining Brown Polia bombycina	Extinct? Last recorded from Rutland in 1995
Pale Tussock Calliteara pudibunda	Common
Pale-shouldered Brocade Lacanobia thalassina	Common
Peach Blossom Thyatira batis	Common
Peacock Moth Macaria notata	Rare, only approx. 12 records, mainly from Rutland.
Pearly Underwing Peridroma saucia	Uncommon migrant
Pebble Hook-tip Drepana falcataria	Fairly common
Pebble Prominent Notodonta ziczac	Common
Peppered Moth Biston betularia	Common
Phoenix Eulithis prunata	Fairly common.
Pine Beauty Panolis flammea	Fairly common.
Pine Carpet Thera firmata	Uncommon/under-recorded? Many records doubtful & few confirmed
Pine Hawk-moth Sphinx pinastri Pinion-spotted Pug Eupithecia insigniata	Uncommon, but increasing and expanding westwards since1995.  Very rare; only 7 post 2000 records.
Pinion-streaked Snout Schrankia costaestrigalis	Uncommon. Under-recorded
Pink-barred Sallow Xanthia togata	Fairly common
Plain Golden Y Autographa jota	Common
Plain Pug Eupithecia simpliciata	Uncommon and only recorded from gardens
Plain Wave Idaea straminata	Uncommon, few confirmed records in recent years.
Poplar Grey Subacronicta megacephala	Common
Poplar Hawk-moth Laothoe populi	Common
Poplar Kitten Furcula bifida	Uncommon
Poplar Lutestring Tethea or	Very rare. Most recent records are from a few Rutland woodland sites.
Powdered Quaker Orthosia gracilis	Fairly common
Pretty Chalk Carpet Melanthia procellata	Very rare; only a handful of records over a widespread area, 2 in last 20yrs.
Privet Hawk-moth Sphinx ligustri	Common apart from in far west of county where it is absent/very rare.
Purple Bar Cosmorhoe ocellata	Fairly common
Purple Clay Diarsia brunnea	Fairly common, especially in Charnwood Forest
Purple Clay Diarsia brunnea Purple Thorn Selenia tetralunaria	Fairly common, especially in Charnwood Forest Common, especially in woodland
Purple Clay Diarsia brunnea Purple Thorn Selenia tetralunaria Puss Moth Cerura vinula	Fairly common, especially in Charnwood Forest Common, especially in woodland Uncommon. Declined dramatically in recent years.
Purple Clay Diarsia brunnea Purple Thorn Selenia tetralunaria Puss Moth Cerura vinula Red Chestnut Cerastis rubricosa	Fairly common, especially in Charnwood Forest Common, especially in woodland Uncommon. Declined dramatically in recent years. Fairly common in woodland, less so in gardens
Purple Clay Diarsia brunnea Purple Thorn Selenia tetralunaria Puss Moth Cerura vinula Red Chestnut Cerastis rubricosa Red Sword-grass Xylena vetusta	Fairly common, especially in Charnwood Forest Common, especially in woodland Uncommon. Declined dramatically in recent years. Fairly common in woodland, less so in gardens Rare. Only 3 modern records: Loughbro', Ulverscroft & Pickworth Gt Wood
Purple Clay Diarsia brunnea Purple Thorn Selenia tetralunaria Puss Moth Cerura vinula Red Chestnut Cerastis rubricosa Red Sword-grass Xylena vetusta Red Twin-spot Carpet Xanthorhoe spadicearia	Fairly common, especially in Charnwood Forest Common, especially in woodland Uncommon. Declined dramatically in recent years. Fairly common in woodland, less so in gardens Rare. Only 3 modern records: Loughbro', Ulverscroft & Pickworth Gt Wood Common
Purple Clay Diarsia brunnea Purple Thorn Selenia tetralunaria Puss Moth Cerura vinula Red Chestnut Cerastis rubricosa Red Sword-grass Xylena vetusta Red Twin-spot Carpet Xanthorhoe spadicearia Red Underwing Catocala nupta	Fairly common, especially in Charnwood Forest Common, especially in woodland Uncommon. Declined dramatically in recent years. Fairly common in woodland, less so in gardens Rare. Only 3 modern records: Loughbro', Ulverscroft & Pickworth Gt Wood Common Fairly common.
Purple Clay Diarsia brunnea Purple Thorn Selenia tetralunaria Puss Moth Cerura vinula Red Chestnut Cerastis rubricosa Red Sword-grass Xylena vetusta Red Twin-spot Carpet Xanthorhoe spadicearia	Fairly common, especially in Charnwood Forest Common, especially in woodland Uncommon. Declined dramatically in recent years. Fairly common in woodland, less so in gardens Rare. Only 3 modern records: Loughbro', Ulverscroft & Pickworth Gt Wood Common
Purple Clay Diarsia brunnea Purple Thorn Selenia tetralunaria Puss Moth Cerura vinula Red Chestnut Cerastis rubricosa Red Sword-grass Xylena vetusta Red Twin-spot Carpet Xanthorhoe spadicearia Red Underwing Catocala nupta Red-belted Clearwing Synanthedon myopaeformis	Fairly common, especially in Charnwood Forest Common, especially in woodland Uncommon. Declined dramatically in recent years. Fairly common in woodland, less so in gardens Rare. Only 3 modern records: Loughbro', Ulverscroft & Pickworth Gt Wood Common Fairly common. Uncommon/under-recorded (can be common at pheromone lures)
Purple Clay Diarsia brunnea Purple Thorn Selenia tetralunaria Puss Moth Cerura vinula Red Chestnut Cerastis rubricosa Red Sword-grass Xylena vetusta Red Twin-spot Carpet Xanthorhoe spadicearia Red Underwing Catocala nupta Red-belted Clearwing Synanthedon myopaeformis Reddish Light Arches Apamea sublustris Red-green Carpet Chloroclysta siterata Red-line Quaker Agrochola lota	Fairly common, especially in Charnwood Forest Common, especially in woodland Uncommon. Declined dramatically in recent years. Fairly common in woodland, less so in gardens Rare. Only 3 modern records: Loughbro', Ulverscroft & Pickworth Gt Wood Common Fairly common. Uncommon/under-recorded (can be common at pheromone lures) Rare. Few post-VCH records: a few from Rutland (mainly) & elsewhere.
Purple Clay Diarsia brunnea Purple Thorn Selenia tetralunaria Puss Moth Cerura vinula Red Chestnut Cerastis rubricosa Red Sword-grass Xylena vetusta Red Twin-spot Carpet Xanthorhoe spadicearia Red Underwing Catocala nupta Red-belted Clearwing Synanthedon myopaeformis Reddish Light Arches Apamea sublustris Red-green Carpet Chloroclysta siterata Red-line Quaker Agrochola lota Red-necked Footman Atolmis rubricollis	Fairly common, especially in Charnwood Forest Common, especially in woodland Uncommon. Declined dramatically in recent years. Fairly common in woodland, less so in gardens Rare. Only 3 modern records: Loughbro', Ulverscroft & Pickworth Gt Wood Common Fairly common. Uncommon/under-recorded (can be common at pheromone lures) Rare. Few post-VCH records: a few from Rutland (mainly) & elsewhere. Common Common Rare, but increasing?
Purple Clay Diarsia brunnea Purple Thorn Selenia tetralunaria Puss Moth Cerura vinula Red Chestnut Cerastis rubricosa Red Sword-grass Xylena vetusta Red Twin-spot Carpet Xanthorhoe spadicearia Red Underwing Catocala nupta Red-belted Clearwing Synanthedon myopaeformis Reddish Light Arches Apamea sublustris Red-green Carpet Chloroclysta siterata Red-line Quaker Agrochola lota Red-necked Footman Atolmis rubricollis Red-tipped Clearwing Synanthedon formicaeformis	Fairly common, especially in Charnwood Forest Common, especially in woodland Uncommon. Declined dramatically in recent years. Fairly common in woodland, less so in gardens Rare. Only 3 modern records: Loughbro', Ulverscroft & Pickworth Gt Wood Common Fairly common. Uncommon/under-recorded (can be common at pheromone lures) Rare. Few post-VCH records: a few from Rutland (mainly) & elsewhere. Common Common Rare, but increasing? Uncommon/under-recorded (post 2005 records using pheromone lures)
Purple Clay Diarsia brunnea Purple Thorn Selenia tetralunaria Puss Moth Cerura vinula Red Chestnut Cerastis rubricosa Red Sword-grass Xylena vetusta Red Twin-spot Carpet Xanthorhoe spadicearia Red Underwing Catocala nupta Red-belted Clearwing Synanthedon myopaeformis Reddish Light Arches Apamea sublustris Red-green Carpet Chloroclysta siterata Red-line Quaker Agrochola lota Red-necked Footman Atolmis rubricollis Red-tipped Clearwing Synanthedon formicaeformis Riband Wave Idaea aversata	Fairly common, especially in Charnwood Forest Common, especially in woodland Uncommon. Declined dramatically in recent years. Fairly common in woodland, less so in gardens Rare. Only 3 modern records: Loughbro', Ulverscroft & Pickworth Gt Wood Common Fairly common. Uncommon/under-recorded (can be common at pheromone lures) Rare. Few post-VCH records: a few from Rutland (mainly) & elsewhere. Common Common Rare, but increasing? Uncommon/under-recorded (post 2005 records using pheromone lures) Common
Purple Clay Diarsia brunnea Purple Thorn Selenia tetralunaria Puss Moth Cerura vinula Red Chestnut Cerastis rubricosa Red Sword-grass Xylena vetusta Red Twin-spot Carpet Xanthorhoe spadicearia Red Underwing Catocala nupta Red-belted Clearwing Synanthedon myopaeformis Reddish Light Arches Apamea sublustris Red-green Carpet Chloroclysta siterata Red-line Quaker Agrochola lota Red-necked Footman Atolmis rubricollis Red-tipped Clearwing Synanthedon formicaeformis Riband Wave Idaea aversata Rivulet Perizoma affinitata	Fairly common, especially in Charnwood Forest Common, especially in woodland Uncommon. Declined dramatically in recent years. Fairly common in woodland, less so in gardens Rare. Only 3 modern records: Loughbro', Ulverscroft & Pickworth Gt Wood Common Fairly common. Uncommon/under-recorded (can be common at pheromone lures) Rare. Few post-VCH records: a few from Rutland (mainly) & elsewhere. Common Common Rare, but increasing? Uncommon/under-recorded (post 2005 records using pheromone lures) Common Common, less so in gardens
Purple Clay Diarsia brunnea Purple Thorn Selenia tetralunaria Puss Moth Cerura vinula Red Chestnut Cerastis rubricosa Red Sword-grass Xylena vetusta Red Twin-spot Carpet Xanthorhoe spadicearia Red Underwing Catocala nupta Red-belted Clearwing Synanthedon myopaeformis Reddish Light Arches Apamea sublustris Red-green Carpet Chloroclysta siterata Red-line Quaker Agrochola lota Red-necked Footman Atolmis rubricollis Red-tipped Clearwing Synanthedon formicaeformis Riband Wave Idaea aversata Rivulet Perizoma affinitata Rosy Footman Miltochrista miniata	Fairly common, especially in Charnwood Forest Common, especially in woodland Uncommon. Declined dramatically in recent years. Fairly common in woodland, less so in gardens Rare. Only 3 modern records: Loughbro', Ulverscroft & Pickworth Gt Wood Common Fairly common. Uncommon/under-recorded (can be common at pheromone lures) Rare. Few post-VCH records: a few from Rutland (mainly) & elsewhere. Common Common Rare, but increasing? Uncommon/under-recorded (post 2005 records using pheromone lures) Common Common, less so in gardens Rare. Only 7 recent records, from 5 sites in Rutland. Expanding?
Purple Clay Diarsia brunnea Purple Thorn Selenia tetralunaria Puss Moth Cerura vinula Red Chestnut Cerastis rubricosa Red Sword-grass Xylena vetusta Red Twin-spot Carpet Xanthorhoe spadicearia Red Underwing Catocala nupta Red-belted Clearwing Synanthedon myopaeformis Reddish Light Arches Apamea sublustris Red-green Carpet Chloroclysta siterata Red-line Quaker Agrochola lota Red-necked Footman Atolmis rubricollis Red-tipped Clearwing Synanthedon formicaeformis Riband Wave Idaea aversata Rivulet Perizoma affinitata Rosy Footman Miltochrista miniata Rosy Marbled Elaphria venustula	Fairly common, especially in Charnwood Forest Common, especially in woodland Uncommon. Declined dramatically in recent years. Fairly common in woodland, less so in gardens Rare. Only 3 modern records: Loughbro', Ulverscroft & Pickworth Gt Wood Common Fairly common. Uncommon/under-recorded (can be common at pheromone lures) Rare. Few post-VCH records: a few from Rutland (mainly) & elsewhere. Common Common Rare, but increasing? Uncommon/under-recorded (post 2005 records using pheromone lures) Common Common, less so in gardens Rare. Only 7 recent records, from 5 sites in Rutland. Expanding? Very rare. Discovered in Swithland Wood in 2014
Purple Clay Diarsia brunnea Purple Thorn Selenia tetralunaria Puss Moth Cerura vinula Red Chestnut Cerastis rubricosa Red Sword-grass Xylena vetusta Red Twin-spot Carpet Xanthorhoe spadicearia Red Underwing Catocala nupta Red-belted Clearwing Synanthedon myopaeformis Reddish Light Arches Apamea sublustris Red-green Carpet Chloroclysta siterata Red-line Quaker Agrochola lota Red-line Quaker Agrochola lota Red-tipped Clearwing Synanthedon formicaeformis Riband Wave Idaea aversata Rivulet Perizoma affinitata Rosy Footman Miltochrista miniata Rosy Marbled Elaphria venustula Rosy Minor Litoligia literosa	Fairly common, especially in Charnwood Forest Common, especially in woodland Uncommon. Declined dramatically in recent years. Fairly common in woodland, less so in gardens Rare. Only 3 modern records: Loughbro', Ulverscroft & Pickworth Gt Wood Common Fairly common. Uncommon/under-recorded (can be common at pheromone lures) Rare. Few post-VCH records: a few from Rutland (mainly) & elsewhere. Common Common Rare, but increasing? Uncommon/under-recorded (post 2005 records using pheromone lures) Common Common, less so in gardens Rare. Only 7 recent records, from 5 sites in Rutland. Expanding? Very rare. Discovered in Swithland Wood in 2014 Fairly common
Purple Clay Diarsia brunnea Purple Thorn Selenia tetralunaria Puss Moth Cerura vinula Red Chestnut Cerastis rubricosa Red Sword-grass Xylena vetusta Red Twin-spot Carpet Xanthorhoe spadicearia Red Underwing Catocala nupta Red-belted Clearwing Synanthedon myopaeformis Reddish Light Arches Apamea sublustris Red-green Carpet Chloroclysta siterata Red-line Quaker Agrochola lota Red-necked Footman Atolmis rubricollis Red-tipped Clearwing Synanthedon formicaeformis Riband Wave Idaea aversata Rivulet Perizoma affinitata Rosy Footman Miltochrista miniata Rosy Marbled Elaphria venustula	Fairly common, especially in Charnwood Forest Common, especially in woodland Uncommon. Declined dramatically in recent years. Fairly common in woodland, less so in gardens Rare. Only 3 modern records: Loughbro', Ulverscroft & Pickworth Gt Wood Common Fairly common. Uncommon/under-recorded (can be common at pheromone lures) Rare. Few post-VCH records: a few from Rutland (mainly) & elsewhere. Common Common Rare, but increasing? Uncommon/under-recorded (post 2005 records using pheromone lures) Common Common, less so in gardens Rare. Only 7 recent records, from 5 sites in Rutland. Expanding? Very rare. Discovered in Swithland Wood in 2014

Ruby Tiger Phragmatobia fuliginosa	Common
Ruddy Carpet Catarhoe rubidata	Very rare. Only 5 post-1970 records, 4 from Rutland
Ruddy Highflier Hydriomena ruberata Rufous Minor Oligia versicolor	Very rare/extinct/status uncertain.
Rufous Minor Oligia Versicolor Rustic Hoplodrina blanda	Fairly common (but few gen-det confirmed ids) Fairly common.
Rustic Shoulder-knot Apamea sordens	Common
Sallow Cirrhia icteritia	Common
Sallow Kitten Furcula furcula	Common
Sandy Carpet Perizoma flavofasciata	Fairly common.
Satellite Eupsilia transversa	Fairly common.
Satin Beauty Deileptenia ribeata	Uncommon & localised.
Satin Wave Idaea subsericeata	Rare & localised. East Rutland only.
Satyr Pug Eupithecia satyrata	Locally common at a few sites in Rutland. Rare elsewhere
Scallop Shell Hydria undulata	Rare. Most frequently recorded at Cloud Wood; very occasional elsewhere.
Scalloped Hazel Odontopera bidentata	Common
Scalloped Hook-tip Falcaria lacertinaria	Locally common
Scalloped Oak Crocallis elinguaria	Common
Scarce Bordered Straw Helicoverpa armigera	Uncommon migrant (common in 2006), plus a few imported larvae records.
Scarce Footman Eilema complana	Fairly common. Dramatically increased in last 10 years.
Scarce Silver-lines Bena bicolorana	Fairly common.
Scarce Tissue Hydria cervinalis	Formerly common, now very rare & only recorded from a few gardens.
Scarce Umber Agriopis aurantiaria	Uncommon and declining
Scarce Vapourer Orgyia recens	Extinct. VCH only.
Scarlet Tiger Callimorpha dominula	Rare. Two known colonies in Rutland but now occasionally elsewhere
Scorched Carpet Ligdia adustata	Uncommon and local, most frequent in Rutland
Scorched Wing Plagodis dolabraria	Fairly common
September Thorn Ennomos erosaria	Uncommon and declining.
Seraphim Lobophora halterata	Fairly common, but local.
Setaceous Hebrew Character Xestia c-nigrum	Common
Shaded Broad-bar Scotopteryx chenopodiata	Common
Shaded Pug Eupithecia subumbrata	Rare. Recent records from a handful of sites in Rutland.
Shark Cucullia umbratica	Fairly common
Sharp-angled Peacock Macaria alternata	Very rare/extinct. VCH (Owston Wood) only
Shears Hada plebeja	Fairly common but local
Short-cloaked Moth Nola cucullatella	Common
Shoulder-stripe Earophila badiata	Common
Shoulder-striped Wainscot Leucania comma	Common
Shuttle-shaped Dart Agrotis puta	Common
Silky Wainscot Chilodes maritima	Common at Rutland Water, only occasional records elsewhere.
Silver Hook Deltote uncula	Extinct. Only known from Narborough Bog; last recorded in 1940.
Silver Y Autographa gamma	Common migrant
Silver-ground Carpet Xanthorhoe montanata	Common
Silver-striped Hawk-moth Hippotion celerio Silvery Arches Polia hepatica	Very rare migrant. Last record in 1963.  Extinct? No recent confirmed record
Single-dotted Wave Idaea dimidiata	Common
Six holted Cleanwing Rembecia ichnoumoniformis	Locally common/under recorded (can be common at phoromone lurge)
Six-belted Clearwing Bembecia ichneumoniformis	Locally common/under-recorded (can be common at pheromone lures)
Six-spot Burnet Zygaena filipendulae	Fairly common. Under-recorded?
Six-spot Burnet Zygaena filipendulae Six-striped Rustic Xestia sexstrigata	Fairly common. Under-recorded? Common
Six-spot Burnet Zygaena filipendulae Six-striped Rustic Xestia sexstrigata Slender Brindle Apamea scolopacina	Fairly common. Under-recorded? Common Fairly common in woodland. Increased since 1980?
Six-spot Burnet Zygaena filipendulae Six-striped Rustic Xestia sexstrigata Slender Brindle Apamea scolopacina Slender Pug Eupithecia tenuiata	Fairly common. Under-recorded? Common Fairly common in woodland. Increased since 1980? Fairly common
Six-spot Burnet Zygaena filipendulae Six-striped Rustic Xestia sexstrigata Slender Brindle Apamea scolopacina Slender Pug Eupithecia tenuiata Sloe Pug Pasiphila chloerata	Fairly common. Under-recorded? Common Fairly common in woodland. Increased since 1980? Fairly common Uncommon.
Six-spot Burnet Zygaena filipendulae Six-striped Rustic Xestia sexstrigata Slender Brindle Apamea scolopacina Slender Pug Eupithecia tenuiata	Fairly common. Under-recorded? Common Fairly common in woodland. Increased since 1980? Fairly common Uncommon. Fairly common.
Six-spot Burnet Zygaena filipendulae Six-striped Rustic Xestia sexstrigata Slender Brindle Apamea scolopacina Slender Pug Eupithecia tenuiata Sloe Pug Pasiphila chloerata Small Angle Shades Euplexia lucipara	Fairly common. Under-recorded? Common Fairly common in woodland. Increased since 1980? Fairly common Uncommon.
Six-spot Burnet Zygaena filipendulae Six-striped Rustic Xestia sexstrigata Slender Brindle Apamea scolopacina Slender Pug Eupithecia tenuiata Sloe Pug Pasiphila chloerata Small Angle Shades Euplexia lucipara Small Argent & Sable Epirrhoe tristata Small Blood-vein Scopula imitaria	Fairly common. Under-recorded?  Common  Fairly common in woodland. Increased since 1980?  Fairly common  Uncommon.  Fairly common.  Only one record which needs investigation.  Fairly common
Six-spot Burnet Zygaena filipendulae Six-striped Rustic Xestia sexstrigata Slender Brindle Apamea scolopacina Slender Pug Eupithecia tenuiata Sloe Pug Pasiphila chloerata Small Angle Shades Euplexia lucipara Small Argent & Sable Epirrhoe tristata	Fairly common. Under-recorded?  Common Fairly common in woodland. Increased since 1980? Fairly common Uncommon. Fairly common. Only one record which needs investigation.
Six-spot Burnet Zygaena filipendulae Six-striped Rustic Xestia sexstrigata Slender Brindle Apamea scolopacina Slender Pug Eupithecia tenuiata Sloe Pug Pasiphila chloerata Small Angle Shades Euplexia lucipara Small Argent & Sable Epirrhoe tristata Small Blood-vein Scopula imitaria Small Brindled Beauty Apocheima hispidaria	Fairly common. Under-recorded?  Common  Fairly common in woodland. Increased since 1980?  Fairly common  Uncommon.  Fairly common.  Only one record which needs investigation.  Fairly common  Rare & localised. Recently rediscovered in Rutland & possibly expanding.
Six-spot Burnet Zygaena filipendulae Six-striped Rustic Xestia sexstrigata Slender Brindle Apamea scolopacina Slender Pug Eupithecia tenuiata Sloe Pug Pasiphila chloerata Small Angle Shades Euplexia lucipara Small Argent & Sable Epirrhoe tristata Small Blood-vein Scopula imitaria Small Brindled Beauty Apocheima hispidaria Small Clouded Brindle Apamea unanimis Small Dotted Buff Photedes minima Small Dusty Wave Idaea seriata	Fairly common. Under-recorded?  Common  Fairly common in woodland. Increased since 1980?  Fairly common  Uncommon.  Fairly common.  Only one record which needs investigation.  Fairly common  Rare & localised. Recently rediscovered in Rutland & possibly expanding.  Uncommon
Six-spot Burnet Zygaena filipendulae Six-striped Rustic Xestia sexstrigata Slender Brindle Apamea scolopacina Slender Pug Eupithecia tenuiata Sloe Pug Pasiphila chloerata Small Angle Shades Euplexia lucipara Small Argent & Sable Epirrhoe tristata Small Blood-vein Scopula imitaria Small Brindled Beauty Apocheima hispidaria Small Clouded Brindle Apamea unanimis Small Dotted Buff Photedes minima Small Dusty Wave Idaea seriata Small Eggar Eriogaster lanestris	Fairly common. Under-recorded?  Common  Fairly common in woodland. Increased since 1980?  Fairly common  Uncommon.  Fairly common.  Only one record which needs investigation.  Fairly common  Rare & localised. Recently rediscovered in Rutland & possibly expanding.  Uncommon  Common  Fairly common.  Very rare/extinct. Few records since VCH. Undiscovered/new colonies?
Six-spot Burnet Zygaena filipendulae Six-striped Rustic Xestia sexstrigata Slender Brindle Apamea scolopacina Slender Pug Eupithecia tenuiata Sloe Pug Pasiphila chloerata Small Angle Shades Euplexia lucipara Small Argent & Sable Epirrhoe tristata Small Blood-vein Scopula imitaria Small Brindled Beauty Apocheima hispidaria Small Clouded Brindle Apamea unanimis Small Dotted Buff Photedes minima Small Dusty Wave Idaea seriata Small Eggar Eriogaster lanestris Small Elephant Hawk-moth Deilephila porcellus	Fairly common. Under-recorded?  Common  Fairly common in woodland. Increased since 1980?  Fairly common  Uncommon.  Fairly common.  Only one record which needs investigation.  Fairly common  Rare & localised. Recently rediscovered in Rutland & possibly expanding.  Uncommon  Common  Fairly common.  Very rare/extinct. Few records since VCH. Undiscovered/new colonies?  Locally common in Charnwood Forest, uncommon/local in rest of county
Six-spot Burnet Zygaena filipendulae Six-striped Rustic Xestia sexstrigata Slender Brindle Apamea scolopacina Slender Pug Eupithecia tenuiata Sloe Pug Pasiphila chloerata Small Angle Shades Euplexia lucipara Small Argent & Sable Epirrhoe tristata Small Blood-vein Scopula imitaria Small Brindled Beauty Apocheima hispidaria Small Clouded Brindle Apamea unanimis Small Dotted Buff Photedes minima Small Dusty Wave Idaea seriata Small Eggar Eriogaster lanestris Small Elephant Hawk-moth Deilephila porcellus Small Emerald Hemistola chrysoprasaria	Fairly common. Under-recorded?  Common  Fairly common in woodland. Increased since 1980?  Fairly common  Uncommon.  Fairly common.  Only one record which needs investigation.  Fairly common  Rare & localised. Recently rediscovered in Rutland & possibly expanding.  Uncommon  Common  Fairly common.  Very rare/extinct. Few records since VCH. Undiscovered/new colonies?  Locally common in Charnwood Forest, uncommon/local in rest of county  Uncommon, but increasing and turning-up in gardens.
Six-spot Burnet Zygaena filipendulae Six-striped Rustic Xestia sexstrigata Slender Brindle Apamea scolopacina Slender Pug Eupithecia tenuiata Sloe Pug Pasiphila chloerata Small Angle Shades Euplexia lucipara Small Argent & Sable Epirrhoe tristata Small Blood-vein Scopula imitaria Small Brindled Beauty Apocheima hispidaria Small Clouded Brindle Apamea unanimis Small Dotted Buff Photedes minima Small Dusty Wave Idaea seriata Small Eggar Eriogaster lanestris Small Elephant Hawk-moth Deilephila porcellus Small Emerald Hemistola chrysoprasaria Small Engrailed Ectropis crepuscularia	Fairly common. Under-recorded?  Common  Fairly common in woodland. Increased since 1980?  Fairly common  Uncommon.  Fairly common.  Only one record which needs investigation.  Fairly common  Rare & localised. Recently rediscovered in Rutland & possibly expanding.  Uncommon  Common  Fairly common.  Very rare/extinct. Few records since VCH. Undiscovered/new colonies?  Locally common in Charnwood Forest, uncommon/local in rest of county  Uncommon, but increasing and turning-up in gardens.  No longer considered to be a distinct species.
Six-spot Burnet Zygaena filipendulae Six-striped Rustic Xestia sexstrigata Slender Brindle Apamea scolopacina Slender Pug Eupithecia tenuiata Sloe Pug Pasiphila chloerata Small Angle Shades Euplexia lucipara Small Argent & Sable Epirrhoe tristata Small Blood-vein Scopula imitaria Small Brindled Beauty Apocheima hispidaria Small Clouded Brindle Apamea unanimis Small Dotted Buff Photedes minima Small Dusty Wave Idaea seriata Small Eggar Eriogaster lanestris Small Elephant Hawk-moth Deilephila porcellus Small Emerald Hemistola chrysoprasaria Small Engrailed Ectropis crepuscularia Small Fan-foot Herminia grisealis	Fairly common. Under-recorded?  Common  Fairly common in woodland. Increased since 1980?  Fairly common  Uncommon.  Fairly common.  Only one record which needs investigation.  Fairly common  Rare & localised. Recently rediscovered in Rutland & possibly expanding.  Uncommon  Common  Fairly common.  Very rare/extinct. Few records since VCH. Undiscovered/new colonies?  Locally common in Charnwood Forest, uncommon/local in rest of county  Uncommon, but increasing and turning-up in gardens.  No longer considered to be a distinct species.  Common
Six-spot Burnet Zygaena filipendulae Six-striped Rustic Xestia sexstrigata Slender Brindle Apamea scolopacina Slender Pug Eupithecia tenuiata Sloe Pug Pasiphila chloerata Small Angle Shades Euplexia lucipara Small Argent & Sable Epirrhoe tristata Small Blood-vein Scopula imitaria Small Brindled Beauty Apocheima hispidaria Small Clouded Brindle Apamea unanimis Small Dotted Buff Photedes minima Small Dusty Wave Idaea seriata Small Eggar Eriogaster lanestris Small Elephant Hawk-moth Deilephila porcellus Small Emerald Hemistola chrysoprasaria Small Engrailed Ectropis crepuscularia Small Fan-foot Herminia grisealis Small Fan-footed Wave Idaea biselata	Fairly common. Under-recorded?  Common  Fairly common in woodland. Increased since 1980?  Fairly common  Uncommon.  Fairly common.  Only one record which needs investigation.  Fairly common  Rare & localised. Recently rediscovered in Rutland & possibly expanding.  Uncommon  Common  Fairly common.  Very rare/extinct. Few records since VCH. Undiscovered/new colonies?  Locally common in Charnwood Forest, uncommon/local in rest of county  Uncommon, but increasing and turning-up in gardens.  No longer considered to be a distinct species.  Common  Common
Six-spot Burnet Zygaena filipendulae Six-striped Rustic Xestia sexstrigata Slender Brindle Apamea scolopacina Slender Pug Eupithecia tenuiata Sloe Pug Pasiphila chloerata Small Angle Shades Euplexia lucipara Small Argent & Sable Epirrhoe tristata Small Blood-vein Scopula imitaria Small Brindled Beauty Apocheima hispidaria Small Clouded Brindle Apamea unanimis Small Dotted Buff Photedes minima Small Dusty Wave Idaea seriata Small Eggar Eriogaster lanestris Small Elephant Hawk-moth Deilephila porcellus Small Emerald Hemistola chrysoprasaria Small Engrailed Ectropis crepuscularia Small Fan-foot Herminia grisealis Small Fan-footed Wave Idaea biselata Small Grass Emerald Chlorissa viridata	Fairly common. Under-recorded?  Common  Fairly common in woodland. Increased since 1980?  Fairly common  Uncommon.  Fairly common.  Only one record which needs investigation.  Fairly common  Rare & localised. Recently rediscovered in Rutland & possibly expanding.  Uncommon  Common  Fairly common.  Very rare/extinct. Few records since VCH. Undiscovered/new colonies?  Locally common in Charnwood Forest, uncommon/local in rest of county  Uncommon, but increasing and turning-up in gardens.  No longer considered to be a distinct species.  Common  Common  Only one record since VCH (Martinshaw Wood in 1980's)
Six-spot Burnet Zygaena filipendulae Six-striped Rustic Xestia sexstrigata Slender Brindle Apamea scolopacina Slender Pug Eupithecia tenuiata Sloe Pug Pasiphila chloerata Small Angle Shades Euplexia lucipara Small Argent & Sable Epirrhoe tristata Small Blood-vein Scopula imitaria Small Brindled Beauty Apocheima hispidaria Small Clouded Brindle Apamea unanimis Small Dotted Buff Photedes minima Small Dusty Wave Idaea seriata Small Eggar Eriogaster lanestris Small Elephant Hawk-moth Deilephila porcellus Small Emerald Hemistola chrysoprasaria Small Engrailed Ectropis crepuscularia Small Fan-foot Herminia grisealis Small Fan-footed Wave Idaea biselata Small Grass Emerald Chlorissa viridata Small Marbled Eublemma parva	Fairly common. Under-recorded?  Common  Fairly common in woodland. Increased since 1980?  Fairly common  Uncommon.  Fairly common.  Only one record which needs investigation.  Fairly common  Rare & localised. Recently rediscovered in Rutland & possibly expanding.  Uncommon  Common  Fairly common.  Very rare/extinct. Few records since VCH. Undiscovered/new colonies?  Locally common in Charnwood Forest, uncommon/local in rest of county  Uncommon, but increasing and turning-up in gardens.  No longer considered to be a distinct species.  Common  Common  Only one record since VCH (Martinshaw Wood in 1980's)  Very rare migrant. Two records: Loughborough (2006) & Littlethorpe (2015)
Six-spot Burnet Zygaena filipendulae Six-striped Rustic Xestia sexstrigata Slender Brindle Apamea scolopacina Slender Pug Eupithecia tenuiata Sloe Pug Pasiphila chloerata Small Angle Shades Euplexia lucipara Small Argent & Sable Epirrhoe tristata Small Blood-vein Scopula imitaria Small Brindled Beauty Apocheima hispidaria Small Clouded Brindle Apamea unanimis Small Dotted Buff Photedes minima Small Dusty Wave Idaea seriata Small Eggar Eriogaster lanestris Small Elephant Hawk-moth Deilephila porcellus Small Emerald Hemistola chrysoprasaria Small Engrailed Ectropis crepuscularia Small Fan-foot Herminia grisealis Small Fan-footed Wave Idaea biselata Small Grass Emerald Chlorissa viridata Small Marbled Eublemma parva Small Mottled Willow Spodoptera exigua	Fairly common. Under-recorded?  Common  Fairly common in woodland. Increased since 1980?  Fairly common  Uncommon.  Fairly common.  Only one record which needs investigation.  Fairly common  Rare & localised. Recently rediscovered in Rutland & possibly expanding.  Uncommon  Common  Fairly common.  Very rare/extinct. Few records since VCH. Undiscovered/new colonies?  Locally common in Charnwood Forest, uncommon/local in rest of county  Uncommon, but increasing and turning-up in gardens.  No longer considered to be a distinct species.  Common  Common  Only one record since VCH (Martinshaw Wood in 1980's)  Very rare migrant. Two records: Loughborough (2006) & Littlethorpe (2015)  Uncommon migrant, but frequently recorded in 2015
Six-spot Burnet Zygaena filipendulae Six-striped Rustic Xestia sexstrigata Slender Brindle Apamea scolopacina Slender Pug Eupithecia tenuiata Sloe Pug Pasiphila chloerata Small Angle Shades Euplexia lucipara Small Argent & Sable Epirrhoe tristata Small Blood-vein Scopula imitaria Small Brindled Beauty Apocheima hispidaria Small Clouded Brindle Apamea unanimis Small Dotted Buff Photedes minima Small Dusty Wave Idaea seriata Small Eggar Eriogaster lanestris Small Elephant Hawk-moth Deilephila porcellus Small Emerald Hemistola chrysoprasaria Small Engrailed Ectropis crepuscularia Small Fan-foot Herminia grisealis Small Fan-footed Wave Idaea biselata Small Grass Emerald Chlorissa viridata Small Marbled Eublemma parva Small Mottled Willow Spodoptera exigua Small Phoenix Ecliptopera silaceata	Fairly common. Under-recorded?  Common  Fairly common in woodland. Increased since 1980?  Fairly common  Uncommon.  Fairly common.  Only one record which needs investigation.  Fairly common  Rare & localised. Recently rediscovered in Rutland & possibly expanding.  Uncommon  Common  Fairly common.  Very rare/extinct. Few records since VCH. Undiscovered/new colonies?  Locally common in Charnwood Forest, uncommon/local in rest of county  Uncommon, but increasing and turning-up in gardens.  No longer considered to be a distinct species.  Common  Common  Only one record since VCH (Martinshaw Wood in 1980's)  Very rare migrant. Two records: Loughborough (2006) & Littlethorpe (2015)  Uncommon migrant, but frequently recorded in 2015  Common
Six-spot Burnet Zygaena filipendulae Six-striped Rustic Xestia sexstrigata Slender Brindle Apamea scolopacina Slender Pug Eupithecia tenuiata Sloe Pug Pasiphila chloerata Small Angle Shades Euplexia lucipara Small Argent & Sable Epirrhoe tristata Small Blood-vein Scopula imitaria Small Brindled Beauty Apocheima hispidaria Small Clouded Brindle Apamea unanimis Small Dotted Buff Photedes minima Small Dusty Wave Idaea seriata Small Eggar Eriogaster lanestris Small Elephant Hawk-moth Deilephila porcellus Small Emerald Hemistola chrysoprasaria Small Engrailed Ectropis crepuscularia Small Fan-foot Herminia grisealis Small Fan-footed Wave Idaea biselata Small Grass Emerald Chlorissa viridata Small Marbled Eublemma parva Small Mottled Willow Spodoptera exigua Small Phoenix Ecliptopera silaceata Small Purple-barred Phytometra viridaria	Fairly common. Under-recorded?  Common  Fairly common in woodland. Increased since 1980?  Fairly common  Uncommon.  Fairly common.  Only one record which needs investigation.  Fairly common  Rare & localised. Recently rediscovered in Rutland & possibly expanding.  Uncommon  Common  Fairly common.  Very rare/extinct. Few records since VCH. Undiscovered/new colonies?  Locally common in Charnwood Forest, uncommon/local in rest of county  Uncommon, but increasing and turning-up in gardens.  No longer considered to be a distinct species.  Common  Common  Only one record since VCH (Martinshaw Wood in 1980's)  Very rare migrant. Two records: Loughborough (2006) & Littlethorpe (2015)  Uncommon migrant, but frequently recorded in 2015  Common  Very rare. Only known colony at Clipsham Quarry/Pickworth Great Wood.
Six-spot Burnet Zygaena filipendulae Six-striped Rustic Xestia sexstrigata Slender Brindle Apamea scolopacina Slender Pug Eupithecia tenuiata Sloe Pug Pasiphila chloerata Small Angle Shades Euplexia lucipara Small Argent & Sable Epirrhoe tristata Small Blood-vein Scopula imitaria Small Brindled Beauty Apocheima hispidaria Small Clouded Brindle Apamea unanimis Small Dotted Buff Photedes minima Small Dusty Wave Idaea seriata Small Eggar Eriogaster lanestris Small Elephant Hawk-moth Deilephila porcellus Small Emerald Hemistola chrysoprasaria Small Engrailed Ectropis crepuscularia Small Fan-foot Herminia grisealis Small Fan-footed Wave Idaea biselata Small Grass Emerald Chlorissa viridata Small Marbled Eublemma parva Small Mottled Willow Spodoptera exigua Small Phoenix Ecliptopera silaceata Small Purple-barred Phytometra viridaria Small Quaker Orthosia cruda	Fairly common. Under-recorded?  Common  Fairly common in woodland. Increased since 1980?  Fairly common  Uncommon.  Fairly common.  Only one record which needs investigation.  Fairly common  Rare & localised. Recently rediscovered in Rutland & possibly expanding.  Uncommon  Common  Fairly common.  Very rare/extinct. Few records since VCH. Undiscovered/new colonies?  Locally common in Charnwood Forest, uncommon/local in rest of county  Uncommon, but increasing and turning-up in gardens.  No longer considered to be a distinct species.  Common  Common  Only one record since VCH (Martinshaw Wood in 1980's)  Very rare migrant. Two records: Loughborough (2006) & Littlethorpe (2015)  Uncommon migrant, but frequently recorded in 2015  Common  Very rare. Only known colony at Clipsham Quarry/Pickworth Great Wood.  Common in suitable habitats.
Six-spot Burnet Zygaena filipendulae Six-striped Rustic Xestia sexstrigata Slender Brindle Apamea scolopacina Slender Pug Eupithecia tenuiata Sloe Pug Pasiphila chloerata Small Angle Shades Euplexia lucipara Small Argent & Sable Epirrhoe tristata Small Blood-vein Scopula imitaria Small Brindled Beauty Apocheima hispidaria Small Clouded Brindle Apamea unanimis Small Dotted Buff Photedes minima Small Dusty Wave Idaea seriata Small Eggar Eriogaster lanestris Small Elephant Hawk-moth Deilephila porcellus Small Emerald Hemistola chrysoprasaria Small Engrailed Ectropis crepuscularia Small Fan-foot Herminia grisealis Small Fan-footed Wave Idaea biselata Small Grass Emerald Chlorissa viridata Small Marbled Eublemma parva Small Mottled Willow Spodoptera exigua Small Phoenix Ecliptopera silaceata Small Purple-barred Phytometra viridaria Small Quaker Orthosia cruda Small Ranunculus Hecatera dysodea	Fairly common. Under-recorded?  Common  Fairly common in woodland. Increased since 1980?  Fairly common  Uncommon.  Fairly common.  Only one record which needs investigation.  Fairly common  Rare & localised. Recently rediscovered in Rutland & possibly expanding.  Uncommon  Common  Fairly common.  Very rare/extinct. Few records since VCH. Undiscovered/new colonies?  Locally common in Charnwood Forest, uncommon/local in rest of county  Uncommon, but increasing and turning-up in gardens.  No longer considered to be a distinct species.  Common  Common  Only one record since VCH (Martinshaw Wood in 1980's)  Very rare migrant. Two records: Loughborough (2006) & Littlethorpe (2015)  Uncommon migrant, but frequently recorded in 2015  Common  Very rare. Only known colony at Clipsham Quarry/Pickworth Great Wood.  Common in suitable habitats.  First recorded in 2006, now locally common (especially on waste ground)
Six-spot Burnet Zygaena filipendulae Six-striped Rustic Xestia sexstrigata Slender Brindle Apamea scolopacina Slender Pug Eupithecia tenuiata Sloe Pug Pasiphila chloerata Small Angle Shades Euplexia lucipara Small Argent & Sable Epirrhoe tristata Small Blood-vein Scopula imitaria Small Brindled Beauty Apocheima hispidaria Small Clouded Brindle Apamea unanimis Small Dotted Buff Photedes minima Small Dusty Wave Idaea seriata Small Eggar Eriogaster lanestris Small Elephant Hawk-moth Deilephila porcellus Small Emerald Hemistola chrysoprasaria Small Engrailed Ectropis crepuscularia Small Fan-foot Herminia grisealis Small Fan-footed Wave Idaea biselata Small Grass Emerald Chlorissa viridata Small Marbled Eublemma parva Small Mottled Willow Spodoptera exigua Small Phoenix Ecliptopera silaceata Small Purple-barred Phytometra viridaria Small Quaker Orthosia cruda Small Ranunculus Hecatera dysodea Small Rivulet Perizoma alchemillata	Fairly common. Under-recorded? Common Fairly common in woodland. Increased since 1980? Fairly common Uncommon. Fairly common. Only one record which needs investigation. Fairly common Rare & localised. Recently rediscovered in Rutland & possibly expanding. Uncommon Common Fairly common. Very rare/extinct. Few records since VCH. Undiscovered/new colonies? Locally common in Charnwood Forest, uncommon/local in rest of county Uncommon, but increasing and turning-up in gardens. No longer considered to be a distinct species. Common Common Only one record since VCH (Martinshaw Wood in 1980's) Very rare migrant. Two records: Loughborough (2006) & Littlethorpe (2015) Uncommon migrant, but frequently recorded in 2015 Common Very rare. Only known colony at Clipsham Quarry/Pickworth Great Wood. Common in suitable habitats. First recorded in 2006, now locally common (especially on waste ground) Common
Six-spot Burnet Zygaena filipendulae Six-striped Rustic Xestia sexstrigata Slender Brindle Apamea scolopacina Slender Pug Eupithecia tenuiata Sloe Pug Pasiphila chloerata Small Angle Shades Euplexia lucipara Small Argent & Sable Epirrhoe tristata Small Blood-vein Scopula imitaria Small Brindled Beauty Apocheima hispidaria Small Clouded Brindle Apamea unanimis Small Dotted Buff Photedes minima Small Dusty Wave Idaea seriata Small Eggar Eriogaster lanestris Small Elephant Hawk-moth Deilephila porcellus Small Enerald Hemistola chrysoprasaria Small Engrailed Ectropis crepuscularia Small Fan-foot Herminia grisealis Small Fan-footed Wave Idaea biselata Small Grass Emerald Chlorissa viridata Small Marbled Eublemma parva Small Mottled Willow Spodoptera exigua Small Phoenix Ecliptopera silaceata Small Purple-barred Phytometra viridaria Small Quaker Orthosia cruda Small Ranunculus Hecatera dysodea Small Rivulet Perizoma alchemillata Small Rufous Coenobia rufa	Fairly common. Under-recorded? Common Fairly common in woodland. Increased since 1980? Fairly common Uncommon. Fairly common. Only one record which needs investigation. Fairly common Rare & localised. Recently rediscovered in Rutland & possibly expanding. Uncommon Common Fairly common. Very rare/extinct. Few records since VCH. Undiscovered/new colonies? Locally common in Charnwood Forest, uncommon/local in rest of county Uncommon, but increasing and turning-up in gardens. No longer considered to be a distinct species. Common Common Only one record since VCH (Martinshaw Wood in 1980's) Very rare migrant. Two records: Loughborough (2006) & Littlethorpe (2015) Uncommon migrant, but frequently recorded in 2015 Common Very rare. Only known colony at Clipsham Quarry/Pickworth Great Wood. Common in suitable habitats. First recorded in 2006, now locally common (especially on waste ground) Common Rare and only recorded from a handful of localities.
Six-spot Burnet Zygaena filipendulae Six-striped Rustic Xestia sexstrigata Slender Brindle Apamea scolopacina Slender Pug Eupithecia tenuiata Sloe Pug Pasiphila chloerata Small Angle Shades Euplexia lucipara Small Argent & Sable Epirrhoe tristata Small Blood-vein Scopula imitaria Small Brindled Beauty Apocheima hispidaria Small Clouded Brindle Apamea unanimis Small Dotted Buff Photedes minima Small Dusty Wave Idaea seriata Small Eggar Eriogaster lanestris Small Elephant Hawk-moth Deilephila porcellus Small Engrailed Ectropis crepuscularia Small Engrailed Ectropis crepuscularia Small Fan-foot Herminia grisealis Small Fan-footed Wave Idaea biselata Small Grass Emerald Chlorissa viridata Small Marbled Eublemma parva Small Mottled Willow Spodoptera exigua Small Phoenix Ecliptopera silaceata Small Purple-barred Phytometra viridaria Small Quaker Orthosia cruda Small Ranunculus Hecatera dysodea Small Rivulet Perizoma alchemillata Small Rufous Coenobia rufa Small Scallop Idaea emarginata	Fairly common in woodland. Increased since 1980? Fairly common in woodland. Increased since 1980? Fairly common Uncommon. Fairly common. Only one record which needs investigation. Fairly common Rare & localised. Recently rediscovered in Rutland & possibly expanding. Uncommon Common Fairly common. Very rare/extinct. Few records since VCH. Undiscovered/new colonies? Locally common in Charnwood Forest, uncommon/local in rest of county Uncommon, but increasing and turning-up in gardens. No longer considered to be a distinct species. Common Common Only one record since VCH (Martinshaw Wood in 1980's) Very rare migrant. Two records: Loughborough (2006) & Littlethorpe (2015) Uncommon migrant, but frequently recorded in 2015 Common Very rare. Only known colony at Clipsham Quarry/Pickworth Great Wood. Common in suitable habitats. First recorded in 2006, now locally common (especially on waste ground) Common Rare and only recorded from a handful of localities. Uncommon & localised. Most records are from eastern half of Rutland.
Six-spot Burnet Zygaena filipendulae Six-striped Rustic Xestia sexstrigata Slender Brindle Apamea scolopacina Slender Pug Eupithecia tenuiata Sloe Pug Pasiphila chloerata Small Angle Shades Euplexia lucipara Small Argent & Sable Epirrhoe tristata Small Blood-vein Scopula imitaria Small Binodevein Scopula imitaria Small Brindled Beauty Apocheima hispidaria Small Clouded Brindle Apamea unanimis Small Dotted Buff Photedes minima Small Dusty Wave Idaea seriata Small Eggar Eriogaster lanestris Small Elephant Hawk-moth Deilephila porcellus Small Engrailed Ectropis crepuscularia Small Engrailed Ectropis crepuscularia Small Fan-foot Herminia grisealis Small Fan-footed Wave Idaea biselata Small Grass Emerald Chlorissa viridata Small Marbled Eublemma parva Small Mottled Willow Spodoptera exigua Small Mottled Willow Spodoptera viridaria Small Phoenix Ecliptopera silaceata Small Purple-barred Phytometra viridaria Small Quaker Orthosia cruda Small Ranunculus Hecatera dysodea Small Rivulet Perizoma alchemillata Small Rufous Coenobia rufa Small Scallop Idaea emarginata Small Square-spot Diarsia rubi	Fairly common. Under-recorded? Common Fairly common in woodland. Increased since 1980? Fairly common Uncommon. Fairly common. Only one record which needs investigation. Fairly common Rare & localised. Recently rediscovered in Rutland & possibly expanding. Uncommon Common Fairly common. Very rare/extinct. Few records since VCH. Undiscovered/new colonies? Locally common in Charnwood Forest, uncommon/local in rest of county Uncommon, but increasing and turning-up in gardens. No longer considered to be a distinct species. Common Common Only one record since VCH (Martinshaw Wood in 1980's) Very rare migrant. Two records: Loughborough (2006) & Littlethorpe (2015) Uncommon migrant, but frequently recorded in 2015 Common Very rare. Only known colony at Clipsham Quarry/Pickworth Great Wood. Common in suitable habitats. First recorded in 2006, now locally common (especially on waste ground) Common Rare and only recorded from a handful of localities. Uncommon, but possibly declining
Six-spot Burnet Zygaena filipendulae Six-striped Rustic Xestia sexstrigata Slender Brindle Apamea scolopacina Slender Pug Eupithecia tenuiata Sloe Pug Pasiphila chloerata Small Angle Shades Euplexia lucipara Small Argent & Sable Epirrhoe tristata Small Blood-vein Scopula imitaria Small Bindled Beauty Apocheima hispidaria Small Clouded Brindle Apamea unanimis Small Dotted Buff Photedes minima Small Dusty Wave Idaea seriata Small Eggar Eriogaster lanestris Small Elephant Hawk-moth Deilephila porcellus Small Emerald Hemistola chrysoprasaria Small Engrailed Ectropis crepuscularia Small Fan-foot Herminia grisealis Small Fan-footed Wave Idaea biselata Small Grass Emerald Chlorissa viridata Small Marbled Eublemma parva Small Mottled Willow Spodoptera exigua Small Mottled Willow Spodoptera exigua Small Phoenix Ecliptopera silaceata Small Purple-barred Phytometra viridaria Small Quaker Orthosia cruda Small Ranunculus Hecatera dysodea Small Rivulet Perizoma alchemillata Small Rufous Coenobia rufa Small Scallop Idaea emarginata Small Square-spot Diarsia rubi Small Wainscot Denticucullus pygmina	Fairly common in woodland. Increased since 1980?  Fairly common  Fairly common  Uncommon.  Only one record which needs investigation.  Fairly common  Rare & localised. Recently rediscovered in Rutland & possibly expanding.  Uncommon  Common  Fairly common.  Very rare/extinct. Few records since VCH. Undiscovered/new colonies?  Locally common in Charnwood Forest, uncommon/local in rest of county  Uncommon, but increasing and turning-up in gardens.  No longer considered to be a distinct species.  Common  Common  Only one record since VCH (Martinshaw Wood in 1980's)  Very rare migrant. Two records: Loughborough (2006) & Littlethorpe (2015)  Uncommon migrant, but frequently recorded in 2015  Common  Very rare. Only known colony at Clipsham Quarry/Pickworth Great Wood.  Common in suitable habitats.  First recorded in 2006, now locally common (especially on waste ground)  Common  Rare and only recorded from a handful of localities.  Uncommon & localised. Most records are from eastern half of Rutland.  Common, but possibly declining  Fairly common
Six-spot Burnet Zygaena filipendulae Six-striped Rustic Xestia sexstrigata Slender Brindle Apamea scolopacina Slender Pug Eupithecia tenuiata Sloe Pug Pasiphila chloerata Small Angle Shades Euplexia lucipara Small Argent & Sable Epirrhoe tristata Small Blood-vein Scopula imitaria Small Binod-vein Scopula imitaria Small Brindled Beauty Apocheima hispidaria Small Clouded Brindle Apamea unanimis Small Dotted Buff Photedes minima Small Dusty Wave Idaea seriata Small Eggar Eriogaster lanestris Small Elephant Hawk-moth Deilephila porcellus Small Emerald Hemistola chrysoprasaria Small Engrailed Ectropis crepuscularia Small Fan-foot Herminia grisealis Small Fan-footed Wave Idaea biselata Small Grass Emerald Chlorissa viridata Small Marbled Eublemma parva Small Mottled Willow Spodoptera exigua Small Mottled Willow Spodoptera exigua Small Phoenix Ecliptopera silaceata Small Purple-barred Phytometra viridaria Small Quaker Orthosia cruda Small Ranunculus Hecatera dysodea Small Rivulet Perizoma alchemillata Small Rivulet Perizoma alchemillata Small Rufous Coenobia rufa Small Scallop Idaea emarginata Small Square-spot Diarsia rubi Small Wainscot Denticucullus pygmina Small Waved Umber Horisme vitalbata	Fairly common. Under-recorded? Common Fairly common in woodland. Increased since 1980? Fairly common Uncommon. Fairly common. Only one record which needs investigation. Fairly common Rare & localised. Recently rediscovered in Rutland & possibly expanding. Uncommon Common Fairly common. Very rare/extinct. Few records since VCH. Undiscovered/new colonies? Locally common in Charrwood Forest, uncommon/local in rest of county Uncommon, but increasing and turning-up in gardens. No longer considered to be a distinct species. Common Common Only one record since VCH (Martinshaw Wood in 1980's) Very rare migrant. Two records: Loughborough (2006) & Littlethorpe (2015) Uncommon migrant, but frequently recorded in 2015 Common Very rare. Only known colony at Clipsham Quarry/Pickworth Great Wood. Common in suitable habitats. First recorded in 2006, now locally common (especially on waste ground) Common Rare and only recorded from a handful of localities. Uncommon & localised. Most records are from eastern half of Rutland. Common, but possibly declining Fairly common Rare, but now appears to be being recorded more frequently in Rutland.
Six-spot Burnet Zygaena filipendulae Six-striped Rustic Xestia sexstrigata Slender Brindle Apamea scolopacina Slender Pug Eupithecia tenuiata Sloe Pug Pasiphila chloerata Small Angle Shades Euplexia lucipara Small Argent & Sable Epirrhoe tristata Small Blood-vein Scopula imitaria Small Binod-vein Scopula imitaria Small Brindled Beauty Apocheima hispidaria Small Clouded Brindle Apamea unanimis Small Dotted Buff Photedes minima Small Dusty Wave Idaea seriata Small Eggar Eriogaster lanestris Small Elephant Hawk-moth Deilephila porcellus Small Emerald Hemistola chrysoprasaria Small Engrailed Ectropis crepuscularia Small Fan-foot Herminia grisealis Small Fan-footed Wave Idaea biselata Small Grass Emerald Chlorissa viridata Small Marbled Eublemma parva Small Mottled Willow Spodoptera exigua Small Phoenix Ecliptopera silaceata Small Purple-barred Phytometra viridaria Small Quaker Orthosia cruda Small Ranunculus Hecatera dysodea Small Rivulet Perizoma alchemillata Small Rufous Coenobia rufa Small Rufous Coenobia rufa Small Scallop Idaea emarginata Small Square-spot Diarsia rubi Small Wainscot Denticucullus pygmina Small Waved Umber Horisme vitalbata Small Waved Umber Horisme vitalbata	Fairly common. Under-recorded? Common Fairly common in woodland. Increased since 1980? Fairly common Uncommon. Fairly common. Only one record which needs investigation. Fairly common Rare & localised. Recently rediscovered in Rutland & possibly expanding. Uncommon Common Fairly common. Very rare/extinct. Few records since VCH. Undiscovered/new colonies? Locally common in Charnwood Forest, uncommon/local in rest of county Uncommon, but increasing and turning-up in gardens. No longer considered to be a distinct species. Common Common Only one record since VCH (Martinshaw Wood in 1980's) Very rare migrant. Two records: Loughborough (2006) & Littlethorpe (2015) Uncommon migrant, but frequently recorded in 2015 Common Very rare. Only known colony at Clipsham Quarry/Pickworth Great Wood. Common in suitable habitats. First recorded in 2006, now locally common (especially on waste ground) Common Rare and only recorded from a handful of localities. Uncommon, but possibly declining Fairly common Rare, but now appears to be being recorded more frequently in Rutland. Becoming uncommon in recent years
Six-spot Burnet Zygaena filipendulae Six-striped Rustic Xestia sexstrigata Slender Brindle Apamea scolopacina Slender Pug Eupithecia tenuiata Sloe Pug Pasiphila chloerata Small Angle Shades Euplexia lucipara Small Argent & Sable Epirrhoe tristata Small Blood-vein Scopula imitaria Small Brindled Beauty Apocheima hispidaria Small Clouded Brindle Apamea unanimis Small Dotted Buff Photedes minima Small Dusty Wave Idaea seriata Small Elephant Hawk-moth Deilephila porcellus Small Elephant Hawk-moth Deilephila porcellus Small Engrailed Ectropis crepuscularia Small Engrailed Ectropis crepuscularia Small Fan-foot Herminia grisealis Small Fan-footed Wave Idaea biselata Small Grass Emerald Chlorissa viridata Small Marbled Eublemma parva Small Mottled Willow Spodoptera exigua Small Phoenix Ecliptopera silaceata Small Purple-barred Phytometra viridaria Small Quaker Orthosia cruda Small Ranunculus Hecatera dysodea Small Rivulet Perizoma alchemillata Small Rufous Coenobia rufa Small Rufous Coenobia rufa Small Scallop Idaea emarginata Small Scallop Idaea emarginata Small Square-spot Diarsia rubi Small Wainscot Denticucullus pygmina Small Waved Umber Horisme vitalbata Small Waved Umber Horisme vitalbata Small White Wave Asthena albulata Small Yellow Underwing Panemeria tenebrata	Fairly common. Under-recorded? Common Fairly common in woodland. Increased since 1980? Fairly common Uncommon. Fairly common. Only one record which needs investigation. Fairly common Rare & localised. Recently rediscovered in Rutland & possibly expanding. Uncommon Common Fairly common. Very rare/extinct. Few records since VCH. Undiscovered/new colonies? Locally common in Charnwood Forest, uncommon/local in rest of county Uncommon, but increasing and turning-up in gardens. No longer considered to be a distinct species. Common Common Only one record since VCH (Martinshaw Wood in 1980's) Very rare migrant. Two records: Loughborough (2006) & Littlethorpe (2015) Uncommon migrant, but frequently recorded in 2015 Common Very rare. Only known colony at Clipsham Quarry/Pickworth Great Wood. Common in suitable habitats. First recorded in 2006, now locally common (especially on waste ground) Common Rare and only recorded from a handful of localities. Uncommon & localised. Most records are from eastern half of Rutland. Common, but possibly declining Fairly common Rare, but now appears to be being recorded more frequently in Rutland. Becoming uncommon in recent years Uncommon and local, possibly declining.
Six-spot Burnet Zygaena filipendulae Six-striped Rustic Xestia sexstrigata Slender Brindle Apamea scolopacina Slender Pug Eupithecia tenuiata Sloe Pug Pasiphila chloerata Small Angle Shades Euplexia lucipara Small Argent & Sable Epirrhoe tristata Small Blood-vein Scopula imitaria Small Bindled Beauty Apocheima hispidaria Small Clouded Brindle Apamea unanimis Small Dotted Buff Photedes minima Small Dusty Wave Idaea seriata Small Eggar Eriogaster lanestris Small Elephant Hawk-moth Deilephila porcellus Small Enerald Hemistola chrysoprasaria Small Engrailed Ectropis crepuscularia Small Fan-foot Herminia grisealis Small Fan-footed Wave Idaea biselata Small Grass Emerald Chlorissa viridata Small Marbled Eublemma parva Small Mottled Willow Spodoptera exigua Small Mottled Willow Spodoptera exigua Small Purple-barred Phytometra viridaria Small Purple-barred Phytometra viridaria Small Ranunculus Hecatera dysodea Small Rivulet Perizoma alchemillata Small Rivulet Perizoma alchemillata Small Rufous Coenobia rufa Small Scallop Idaea emarginata Small Square-spot Diarsia rubi Small Wainscot Denticucullus pygmina Small Waved Umber Horisme vitalbata Small Waved Umber Horisme vitalbata	Fairly common. Under-recorded? Common Fairly common in woodland. Increased since 1980? Fairly common Uncommon. Fairly common. Only one record which needs investigation. Fairly common Rare & localised. Recently rediscovered in Rutland & possibly expanding. Uncommon Common Fairly common. Very rare/extinct. Few records since VCH. Undiscovered/new colonies? Locally common in Charnwood Forest, uncommon/local in rest of county Uncommon, but increasing and turning-up in gardens. No longer considered to be a distinct species. Common Common Only one record since VCH (Martinshaw Wood in 1980's) Very rare migrant. Two records: Loughborough (2006) & Littlethorpe (2015) Uncommon migrant, but frequently recorded in 2015 Common Very rare. Only known colony at Clipsham Quarry/Pickworth Great Wood. Common in suitable habitats. First recorded in 2006, now locally common (especially on waste ground) Common Rare and only recorded from a handful of localities. Uncommon, but possibly declining Fairly common Rare, but now appears to be being recorded more frequently in Rutland. Becoming uncommon in recent years

Snout Hypena proboscidalis	Common
Southern Wainscot Mythimna straminea	Locally common in reedbeds.
Speckled Yellow Pseudopanthera macularia	Very rare/extinct (a couple of recent unconfirmed records).
Spectacle Abrostola tripartita	Common
Spinach Eulithis mellinata	Common
Sprawler Asteroscopus sphinx	Fairly common/under-recorded
Spring Usher Agriopis leucophaearia	Fairly common. Under-recorded?  Common, particularly in gardens. Increased in last 20 years.
Spruce Carpet Thera britannica Square Spot Paradarisa consonaria	Very rare/extinct. Only a couple of 1980's records and one in 2005.
Square-spot Rustic Xestia xanthographa	Common
Square-spotted Clay Xestia stigmatica	Locally common in Rutland, very rare or absent elsewhere
Stout Dart Spaelotis ravida	Declined and extinct (similar situation nationally)
Straw Dot Rivula sericealis	Common.
Straw Underwing Thalpophila matura	Common.
Streak Chesias legatella	Uncommon & local. Occasionally found in good numbers.
Streamer Anticlea derivata Striped Hawk-moth Hyles lineata	Common  Very rare migrant. Only 5 post-VCH records.
Striped Twin-spot Carpet Coenotephria salicata	VCH (Owston Wood) & recent records from RITs only. Very rare
Suspected Parastichtis suspecta	Uncommon & localised. Most frequently recorded in Charnwood Forest.
Svensson's Copper Underwing Amphipyra berbera	Uncommon/under-recorded?
Swallow Prominent Pheosia tremula	Common
Swallow-tailed Moth Ourapteryx sambucaria	Common
Sword-grass Xylena exsoleta	Probably extinct. A few recent records require investigation.
Sycamore Acronicta aceris  Taway Marbled Minor Oligia latrungula	Locally common. Major increase since 1990, but still commonest in east.  Fairly common (but few gen-det confirmed ids)
Tawny Marbled Minor Oligia latruncula Tawny Pinion Lithophane semibrunnea	Uncommon (but few gen-det confirmed ids)  Uncommon & local, but increased in last 10 years.
Tawny Shears Hadena perplexa	Uncommon but local in Rutland, rare in west.
Tawny Speckled Pug Eupithecia icterata	Fairly common.
Tawny-barred Angle Macaria liturata	Fairly common in wooded areas
Tissue Triphosa dubitata	Uncommon and fairly local.
Toadflax Pug Eupithecia linariata	Rare. Only recorded from a handful of gardens in recent years. (
Treble Brown Spot Idaea trigeminata	Rare, but increasing
Treble Lines Charanyca trigrammica Treble-bar Aplocera plagiata	Common in Rutland, uncommon elsewhere Fairly common
Triple-spotted Clay Xestia ditrapezium	Rare/uncommon & declining? Mainly recorded from Charnwood Forest.
Triple-spotted Pug Eupithecia trisignaria	Uncommon, particularly in recent years. Mainly recorded from west of county
True Lover's Knot Lycophotia porphyrea	Locally common in gardens and in Charnwood Forest.
T : 14 () 4 ()	
Turnip Moth Agrotis segetum	Common
Twin-spot Carpet Mesotype didymata	Fairly common
Twin-spot Carpet Mesotype didymata Twin-spotted Quaker Anorthoa munda	Fairly common Fairly common in suitable habitats.
Twin-spot Carpet Mesotype didymata Twin-spotted Quaker Anorthoa munda Twin-spotted Wainscot Lenisa geminipuncta	Fairly common Fairly common in suitable habitats. Uncommon. Mainly recorded from Rutland Water, but occurs elsewhere
Twin-spot Carpet Mesotype didymata Twin-spotted Quaker Anorthoa munda Twin-spotted Wainscot Lenisa geminipuncta Uncertain Hoplodrina octogenaria	Fairly common Fairly common in suitable habitats. Uncommon. Mainly recorded from Rutland Water, but occurs elsewhere Common.
Twin-spot Carpet Mesotype didymata Twin-spotted Quaker Anorthoa munda Twin-spotted Wainscot Lenisa geminipuncta	Fairly common Fairly common in suitable habitats. Uncommon. Mainly recorded from Rutland Water, but occurs elsewhere
Twin-spot Carpet Mesotype didymata Twin-spotted Quaker Anorthoa munda Twin-spotted Wainscot Lenisa geminipuncta Uncertain Hoplodrina octogenaria Valerian Pug Eupithecia valerianata Vapourer Orgyia antiqua Varied Coronet Hadena compta	Fairly common Fairly common in suitable habitats. Uncommon. Mainly recorded from Rutland Water, but occurs elsewhere Common. Only three reliable records, from two woodland sites in east of county. Common especially in urban areas Now fairly common in gardens
Twin-spot Carpet Mesotype didymata Twin-spotted Quaker Anorthoa munda Twin-spotted Wainscot Lenisa geminipuncta Uncertain Hoplodrina octogenaria Valerian Pug Eupithecia valerianata Vapourer Orgyia antiqua Varied Coronet Hadena compta Vestal Rhodometra sacraria	Fairly common Fairly common in suitable habitats. Uncommon. Mainly recorded from Rutland Water, but occurs elsewhere Common. Only three reliable records, from two woodland sites in east of county. Common especially in urban areas Now fairly common in gardens Formerly an uncommon migrant, now seen more frequently.
Twin-spot Carpet Mesotype didymata Twin-spotted Quaker Anorthoa munda Twin-spotted Wainscot Lenisa geminipuncta Uncertain Hoplodrina octogenaria Valerian Pug Eupithecia valerianata Vapourer Orgyia antiqua Varied Coronet Hadena compta Vestal Rhodometra sacraria Vine's Rustic Hoplodrina ambigua	Fairly common Fairly common in suitable habitats. Uncommon. Mainly recorded from Rutland Water, but occurs elsewhere Common. Only three reliable records, from two woodland sites in east of county. Common especially in urban areas Now fairly common in gardens Formerly an uncommon migrant, now seen more frequently. Common, especially in eastern half of county
Twin-spot Carpet Mesotype didymata Twin-spotted Quaker Anorthoa munda Twin-spotted Wainscot Lenisa geminipuncta Uncertain Hoplodrina octogenaria Valerian Pug Eupithecia valerianata Vapourer Orgyia antiqua Varied Coronet Hadena compta Vestal Rhodometra sacraria Vine's Rustic Hoplodrina ambigua V-Moth Macaria wauaria	Fairly common Fairly common in suitable habitats. Uncommon. Mainly recorded from Rutland Water, but occurs elsewhere Common. Only three reliable records, from two woodland sites in east of county. Common especially in urban areas Now fairly common in gardens Formerly an uncommon migrant, now seen more frequently. Common, especially in eastern half of county Very Rare. Declined dramatically. Now only know from Clipsham Quarry
Twin-spot Carpet Mesotype didymata Twin-spotted Quaker Anorthoa munda Twin-spotted Wainscot Lenisa geminipuncta Uncertain Hoplodrina octogenaria Valerian Pug Eupithecia valerianata Vapourer Orgyia antiqua Varied Coronet Hadena compta Vestal Rhodometra sacraria Vine's Rustic Hoplodrina ambigua V-Moth Macaria wauaria V-Pug Chloroclystis v-ata	Fairly common Fairly common in suitable habitats. Uncommon. Mainly recorded from Rutland Water, but occurs elsewhere Common. Only three reliable records, from two woodland sites in east of county. Common especially in urban areas Now fairly common in gardens Formerly an uncommon migrant, now seen more frequently. Common, especially in eastern half of county Very Rare. Declined dramatically. Now only know from Clipsham Quarry Common.
Twin-spot Carpet Mesotype didymata Twin-spotted Quaker Anorthoa munda Twin-spotted Wainscot Lenisa geminipuncta Uncertain Hoplodrina octogenaria Valerian Pug Eupithecia valerianata Vapourer Orgyia antiqua Varied Coronet Hadena compta Vestal Rhodometra sacraria Vine's Rustic Hoplodrina ambigua V-Moth Macaria wauaria V-Pug Chloroclystis v-ata Water Carpet Lampropteryx suffumata	Fairly common Fairly common in suitable habitats. Uncommon. Mainly recorded from Rutland Water, but occurs elsewhere Common. Only three reliable records, from two woodland sites in east of county. Common especially in urban areas Now fairly common in gardens Formerly an uncommon migrant, now seen more frequently. Common, especially in eastern half of county Very Rare. Declined dramatically. Now only know from Clipsham Quarry Common. Fairly common
Twin-spot Carpet Mesotype didymata Twin-spotted Quaker Anorthoa munda Twin-spotted Wainscot Lenisa geminipuncta Uncertain Hoplodrina octogenaria Valerian Pug Eupithecia valerianata Vapourer Orgyia antiqua Varied Coronet Hadena compta Vestal Rhodometra sacraria Vine's Rustic Hoplodrina ambigua V-Moth Macaria wauaria V-Pug Chloroclystis v-ata	Fairly common Fairly common in suitable habitats. Uncommon. Mainly recorded from Rutland Water, but occurs elsewhere Common. Only three reliable records, from two woodland sites in east of county. Common especially in urban areas Now fairly common in gardens Formerly an uncommon migrant, now seen more frequently. Common, especially in eastern half of county Very Rare. Declined dramatically. Now only know from Clipsham Quarry Common. Fairly common Extinct. VCH & a couple of other records which need investigating Very rare. Only six records, all recent. Increasing?
Twin-spot Carpet Mesotype didymata Twin-spotted Quaker Anorthoa munda Twin-spotted Wainscot Lenisa geminipuncta Uncertain Hoplodrina octogenaria Valerian Pug Eupithecia valerianata Vapourer Orgyia antiqua Varied Coronet Hadena compta Vestal Rhodometra sacraria Vine's Rustic Hoplodrina ambigua V-Moth Macaria wauaria V-Pug Chloroclystis v-ata Water Carpet Lampropteryx suffumata Water Ermine Spilosoma urticae Waved Black Parascotia fuliginaria Waved Umber Menophra abruptaria	Fairly common Fairly common in suitable habitats. Uncommon. Mainly recorded from Rutland Water, but occurs elsewhere Common. Only three reliable records, from two woodland sites in east of county. Common especially in urban areas Now fairly common in gardens Formerly an uncommon migrant, now seen more frequently. Common, especially in eastern half of county Very Rare. Declined dramatically. Now only know from Clipsham Quarry Common. Fairly common Extinct. VCH & a couple of other records which need investigating Very rare. Only six records, all recent. Increasing? Common
Twin-spot Carpet Mesotype didymata Twin-spotted Quaker Anorthoa munda Twin-spotted Wainscot Lenisa geminipuncta Uncertain Hoplodrina octogenaria Valerian Pug Eupithecia valerianata Vapourer Orgyia antiqua Varied Coronet Hadena compta Vestal Rhodometra sacraria Vine's Rustic Hoplodrina ambigua V-Moth Macaria wauaria V-Pug Chloroclystis v-ata Water Carpet Lampropteryx suffumata Water Ermine Spilosoma urticae Waved Black Parascotia fuliginaria Waved Umber Menophra abruptaria Webb's Wainscot Globia sparganii	Fairly common Fairly common in suitable habitats. Uncommon. Mainly recorded from Rutland Water, but occurs elsewhere Common. Only three reliable records, from two woodland sites in east of county. Common especially in urban areas Now fairly common in gardens Formerly an uncommon migrant, now seen more frequently. Common, especially in eastern half of county Very Rare. Declined dramatically. Now only know from Clipsham Quarry Common. Fairly common Extinct. VCH & a couple of other records which need investigating Very rare. Only six records, all recent. Increasing? Common Very rare. Only a handful of recent records
Twin-spot Carpet Mesotype didymata Twin-spotted Quaker Anorthoa munda Twin-spotted Wainscot Lenisa geminipuncta Uncertain Hoplodrina octogenaria Valerian Pug Eupithecia valerianata Vapourer Orgyia antiqua Varied Coronet Hadena compta Vestal Rhodometra sacraria Vine's Rustic Hoplodrina ambigua V-Moth Macaria wauaria V-Pug Chloroclystis v-ata Water Carpet Lampropteryx suffumata Water Ermine Spilosoma urticae Waved Black Parascotia fuliginaria Waved Umber Menophra abruptaria Webb's Wainscot Globia sparganii White Ermine Spilosoma lubricipeda	Fairly common Fairly common in suitable habitats. Uncommon. Mainly recorded from Rutland Water, but occurs elsewhere Common. Only three reliable records, from two woodland sites in east of county. Common especially in urban areas Now fairly common in gardens Formerly an uncommon migrant, now seen more frequently. Common, especially in eastern half of county Very Rare. Declined dramatically. Now only know from Clipsham Quarry Common. Fairly common Extinct. VCH & a couple of other records which need investigating Very rare. Only six records, all recent. Increasing? Common Very rare. Only a handful of recent records Common
Twin-spot Carpet Mesotype didymata Twin-spotted Quaker Anorthoa munda Twin-spotted Wainscot Lenisa geminipuncta Uncertain Hoplodrina octogenaria Valerian Pug Eupithecia valerianata Vapourer Orgyia antiqua Varied Coronet Hadena compta Vestal Rhodometra sacraria Vine's Rustic Hoplodrina ambigua V-Moth Macaria wauaria V-Pug Chloroclystis v-ata Water Carpet Lampropteryx suffumata Water Ermine Spilosoma urticae Waved Black Parascotia fuliginaria Waved Umber Menophra abruptaria Webb's Wainscot Globia sparganii White Ermine Spilosoma lubricipeda White Satin Moth Leucoma salicis	Fairly common Fairly common in suitable habitats.  Uncommon. Mainly recorded from Rutland Water, but occurs elsewhere Common. Only three reliable records, from two woodland sites in east of county. Common especially in urban areas Now fairly common in gardens Formerly an uncommon migrant, now seen more frequently. Common, especially in eastern half of county Very Rare. Declined dramatically. Now only know from Clipsham Quarry Common. Fairly common Extinct. VCH & a couple of other records which need investigating Very rare. Only six records, all recent. Increasing? Common Very rare. Only a handful of recent records Common Fairly common
Twin-spot Carpet Mesotype didymata Twin-spotted Quaker Anorthoa munda Twin-spotted Wainscot Lenisa geminipuncta Uncertain Hoplodrina octogenaria Valerian Pug Eupithecia valerianata Vapourer Orgyia antiqua Varied Coronet Hadena compta Vestal Rhodometra sacraria Vine's Rustic Hoplodrina ambigua V-Moth Macaria wauaria V-Pug Chloroclystis v-ata Water Carpet Lampropteryx suffumata Water Ermine Spilosoma urticae Waved Black Parascotia fuliginaria Webb's Wainscot Globia sparganii White Ermine Spilosoma lubricipeda White Satin Moth Leucoma salicis White-line Dart Euxoa tritici	Fairly common Fairly common in suitable habitats.  Uncommon. Mainly recorded from Rutland Water, but occurs elsewhere Common. Only three reliable records, from two woodland sites in east of county. Common especially in urban areas Now fairly common in gardens Formerly an uncommon migrant, now seen more frequently. Common, especially in eastern half of county Very Rare. Declined dramatically. Now only know from Clipsham Quarry Common. Fairly common Extinct. VCH & a couple of other records which need investigating Very rare. Only six records, all recent. Increasing? Common Very rare. Only a handful of recent records Common Fairly common Fairly common Fairly common
Twin-spot Carpet Mesotype didymata Twin-spotted Quaker Anorthoa munda Twin-spotted Wainscot Lenisa geminipuncta Uncertain Hoplodrina octogenaria Valerian Pug Eupithecia valerianata Vapourer Orgyia antiqua Varied Coronet Hadena compta Vestal Rhodometra sacraria Vine's Rustic Hoplodrina ambigua V-Moth Macaria wauaria V-Pug Chloroclystis v-ata Water Carpet Lampropteryx suffumata Water Ermine Spilosoma urticae Waved Black Parascotia fuliginaria Waved Umber Menophra abruptaria Webb's Wainscot Globia sparganii White Ermine Spilosoma lubricipeda White Satin Moth Leucoma salicis White-line Dart Euxoa tritici White-marked Cerastis leucographa	Fairly common Fairly common in suitable habitats.  Uncommon. Mainly recorded from Rutland Water, but occurs elsewhere Common. Only three reliable records, from two woodland sites in east of county. Common especially in urban areas Now fairly common in gardens Formerly an uncommon migrant, now seen more frequently. Common, especially in eastern half of county Very Rare. Declined dramatically. Now only know from Clipsham Quarry Common. Fairly common Extinct. VCH & a couple of other records which need investigating Very rare. Only six records, all recent. Increasing? Common Very rare. Only a handful of recent records Common Fairly common
Twin-spot Carpet Mesotype didymata Twin-spotted Quaker Anorthoa munda Twin-spotted Wainscot Lenisa geminipuncta Uncertain Hoplodrina octogenaria Valerian Pug Eupithecia valerianata Vapourer Orgyia antiqua Varied Coronet Hadena compta Vestal Rhodometra sacraria Vine's Rustic Hoplodrina ambigua V-Moth Macaria wauaria V-Pug Chloroclystis v-ata Water Carpet Lampropteryx suffumata Water Ermine Spilosoma urticae Waved Black Parascotia fuliginaria Webb's Wainscot Globia sparganii White Ermine Spilosoma lubricipeda White Satin Moth Leucoma salicis White-line Dart Euxoa tritici White-marked Cerastis leucographa White-point Mythimna albipuncta	Fairly common Fairly common in suitable habitats.  Uncommon. Mainly recorded from Rutland Water, but occurs elsewhere Common. Only three reliable records, from two woodland sites in east of county. Common especially in urban areas Now fairly common in gardens Formerly an uncommon migrant, now seen more frequently. Common, especially in eastern half of county Very Rare. Declined dramatically. Now only know from Clipsham Quarry Common. Fairly common Extinct. VCH & a couple of other records which need investigating Very rare. Only six records, all recent. Increasing? Common Very rare. Only a handful of recent records Common Fairly common Fairly common Fairly rare Rare & local, restricted to the east of County Dramatic increase since mid-1980's, now common, less so in gardens Very rare migrant. Only a handful of recent records. (
Twin-spot Carpet Mesotype didymata Twin-spotted Quaker Anorthoa munda Twin-spotted Wainscot Lenisa geminipuncta Uncertain Hoplodrina octogenaria Valerian Pug Eupithecia valerianata Vapourer Orgyia antiqua Varied Coronet Hadena compta Vestal Rhodometra sacraria Vine's Rustic Hoplodrina ambigua V-Moth Macaria wauaria V-Pug Chloroclystis v-ata Water Carpet Lampropteryx suffumata Water Ermine Spilosoma urticae Waved Black Parascotia fuliginaria Waved Umber Menophra abruptaria Webb's Wainscot Globia sparganii White Ermine Spilosoma lubricipeda White Satin Moth Leucoma salicis White-line Dart Euxoa tritici White-marked Cerastis leucographa White-point Mythimna albipuncta White-spotted Pinion Cosmia diffinis	Fairly common Fairly common in suitable habitats.  Uncommon. Mainly recorded from Rutland Water, but occurs elsewhere Common. Only three reliable records, from two woodland sites in east of county. Common especially in urban areas Now fairly common in gardens Formerly an uncommon migrant, now seen more frequently. Common, especially in eastern half of county Very Rare. Declined dramatically. Now only know from Clipsham Quarry Common. Fairly common Extinct. VCH & a couple of other records which need investigating Very rare. Only six records, all recent. Increasing? Common Very rare. Only a handful of recent records Common Fairly common Fairly common Fairly rare Rare & local, restricted to the east of County Dramatic increase since mid-1980's, now common, less so in gardens Very rare migrant. Only a handful of recent records. ( Extinct? Only two, possibly doubtful, records since 1970.
Twin-spot Carpet Mesotype didymata Twin-spotted Quaker Anorthoa munda Twin-spotted Wainscot Lenisa geminipuncta Uncertain Hoplodrina octogenaria Valerian Pug Eupithecia valerianata Vapourer Orgyia antiqua Varied Coronet Hadena compta Vestal Rhodometra sacraria Vine's Rustic Hoplodrina ambigua V-Moth Macaria wauaria V-Pug Chloroclystis v-ata Water Carpet Lampropteryx suffumata Water Ermine Spilosoma urticae Waved Black Parascotia fuliginaria Waved Umber Menophra abruptaria Webb's Wainscot Globia sparganii White Ermine Spilosoma lubricipeda White Satin Moth Leucoma salicis White-line Dart Euxoa tritici White-marked Cerastis leucographa White-point Mythimna albipuncta White-spotted Pinion Cosmia diffinis White-spotted Pug Eupithecia tripunctaria	Fairly common Fairly common in suitable habitats.  Uncommon. Mainly recorded from Rutland Water, but occurs elsewhere Common. Only three reliable records, from two woodland sites in east of county. Common especially in urban areas Now fairly common in gardens Formerly an uncommon migrant, now seen more frequently. Common, especially in eastern half of county Very Rare. Declined dramatically. Now only know from Clipsham Quarry Common. Fairly common Extinct. VCH & a couple of other records which need investigating Very rare. Only six records, all recent. Increasing? Common Very rare. Only a handful of recent records Common Fairly common Fairly common Fairly rare Rare & local, restricted to the east of County Dramatic increase since mid-1980's, now common, less so in gardens Very rare migrant. Only a handful of recent records. ( Extinct? Only two, possibly doubtful, records since 1970.
Twin-spot Carpet Mesotype didymata Twin-spotted Quaker Anorthoa munda Twin-spotted Wainscot Lenisa geminipuncta Uncertain Hoplodrina octogenaria Valerian Pug Eupithecia valerianata Vapourer Orgyia antiqua Varied Coronet Hadena compta Vestal Rhodometra sacraria Vine's Rustic Hoplodrina ambigua V-Moth Macaria wauaria V-Pug Chloroclystis v-ata Water Carpet Lampropteryx suffumata Water Ermine Spilosoma urticae Waved Black Parascotia fuliginaria Waved Umber Menophra abruptaria Webb's Wainscot Globia sparganii White Ermine Spilosoma lubricipeda White Satin Moth Leucoma salicis White-line Dart Euxoa tritici White-marked Cerastis leucographa White-point Mythimna albipuncta White-spotted Pinion Cosmia diffinis White-spotted Pug Eupithecia tripunctaria Willow Beauty Peribatodes rhomboidaria	Fairly common Fairly common in suitable habitats.  Uncommon. Mainly recorded from Rutland Water, but occurs elsewhere Common. Only three reliable records, from two woodland sites in east of county. Common especially in urban areas Now fairly common in gardens Formerly an uncommon migrant, now seen more frequently. Common, especially in eastern half of county Very Rare. Declined dramatically. Now only know from Clipsham Quarry Common. Fairly common Extinct. VCH & a couple of other records which need investigating Very rare. Only six records, all recent. Increasing? Common Very rare. Only a handful of recent records Common Fairly common Fairly common Fairly rare Rare & local, restricted to the east of County Dramatic increase since mid-1980's, now common, less so in gardens Very rare migrant. Only a handful of recent records. ( Extinct? Only two, possibly doubtful, records since 1970. Common
Twin-spot Carpet Mesotype didymata Twin-spotted Quaker Anorthoa munda Twin-spotted Wainscot Lenisa geminipuncta Uncertain Hoplodrina octogenaria Valerian Pug Eupithecia valerianata Vapourer Orgyia antiqua Varied Coronet Hadena compta Vestal Rhodometra sacraria Vine's Rustic Hoplodrina ambigua V-Moth Macaria wauaria V-Pug Chloroclystis v-ata Water Carpet Lampropteryx suffumata Water Ermine Spilosoma urticae Waved Black Parascotia fuliginaria Waved Umber Menophra abruptaria Webb's Wainscot Globia sparganii White Ermine Spilosoma lubricipeda White Satin Moth Leucoma salicis White-line Dart Euxoa tritici White-marked Cerastis leucographa White-point Mythimna albipuncta White-spotted Pinion Cosmia diffinis White-spotted Pug Eupithecia tripunctaria Willow Beauty Peribatodes rhomboidaria Winter Moth Operophtera brumata	Fairly common Fairly common in suitable habitats.  Uncommon. Mainly recorded from Rutland Water, but occurs elsewhere Common. Only three reliable records, from two woodland sites in east of county. Common especially in urban areas Now fairly common in gardens Formerly an uncommon migrant, now seen more frequently. Common, especially in eastern half of county Very Rare. Declined dramatically. Now only know from Clipsham Quarry Common. Fairly common Extinct. VCH & a couple of other records which need investigating Very rare. Only six records, all recent. Increasing? Common Very rare. Only a handful of recent records Common Fairly common Fairly common Fairly common Fairly rare Rare & local, restricted to the east of County Dramatic increase since mid-1980's, now common, less so in gardens Very rare migrant. Only a handful of recent records. ( Extinct? Only two, possibly doubtful, records since 1970. Common Common
Twin-spot Carpet Mesotype didymata Twin-spotted Quaker Anorthoa munda Twin-spotted Wainscot Lenisa geminipuncta Uncertain Hoplodrina octogenaria Valerian Pug Eupithecia valerianata Vapourer Orgyia antiqua Varied Coronet Hadena compta Vestal Rhodometra sacraria Vine's Rustic Hoplodrina ambigua V-Moth Macaria wauaria V-Pug Chloroclystis v-ata Water Carpet Lampropteryx suffumata Water Ermine Spilosoma urticae Waved Black Parascotia fuliginaria Waved Umber Menophra abruptaria Webb's Wainscot Globia sparganii White Ermine Spilosoma lubricipeda White Satin Moth Leucoma salicis White-line Dart Euxoa tritici White-marked Cerastis leucographa White-point Mythimna albipuncta White-spotted Pinion Cosmia diffinis White-spotted Pug Eupithecia tripunctaria Willow Beauty Peribatodes rhomboidaria Winter Moth Operophtera brumata Wood Carpet Epirrhoe rivata	Fairly common Fairly common in suitable habitats. Uncommon. Mainly recorded from Rutland Water, but occurs elsewhere Common. Only three reliable records, from two woodland sites in east of county. Common especially in urban areas Now fairly common in gardens Formerly an uncommon migrant, now seen more frequently. Common, especially in eastern half of county Very Rare. Declined dramatically. Now only know from Clipsham Quarry Common. Fairly common Extinct. VCH & a couple of other records which need investigating Very rare. Only six records, all recent. Increasing? Common Very rare. Only a handful of recent records Common Fairly common Fairly common Fairly rare Rare & local, restricted to the east of County Dramatic increase since mid-1980's, now common, less so in gardens Very rare migrant. Only a handful of recent records. ( Extinct? Only two, possibly doubtful, records since 1970. Common Common Common Very rare/status uncertain - all recent records need checking/verification.
Twin-spot Carpet Mesotype didymata Twin-spotted Quaker Anorthoa munda Twin-spotted Wainscot Lenisa geminipuncta Uncertain Hoplodrina octogenaria Valerian Pug Eupithecia valerianata Vapourer Orgyia antiqua Varied Coronet Hadena compta Vestal Rhodometra sacraria Vine's Rustic Hoplodrina ambigua V-Moth Macaria wauaria V-Pug Chloroclystis v-ata Water Carpet Lampropteryx suffumata Water Ermine Spilosoma urticae Waved Black Parascotia fuliginaria Waved Umber Menophra abruptaria Webb's Wainscot Globia sparganii White Ermine Spilosoma lubricipeda White Satin Moth Leucoma salicis White-line Dart Euxoa tritici White-marked Cerastis leucographa White-point Mythimna albipuncta White-spotted Pinion Cosmia diffinis White-spotted Pug Eupithecia tripunctaria Willow Beauty Peribatodes rhomboidaria Winter Moth Operophtera brumata	Fairly common Fairly common in suitable habitats.  Uncommon. Mainly recorded from Rutland Water, but occurs elsewhere Common. Only three reliable records, from two woodland sites in east of county. Common especially in urban areas Now fairly common in gardens Formerly an uncommon migrant, now seen more frequently. Common, especially in eastern half of county Very Rare. Declined dramatically. Now only know from Clipsham Quarry Common. Fairly common Extinct. VCH & a couple of other records which need investigating Very rare. Only six records, all recent. Increasing? Common Very rare. Only a handful of recent records Common Fairly common Fairly common Fairly common Fairly rare Rare & local, restricted to the east of County Dramatic increase since mid-1980's, now common, less so in gardens Very rare migrant. Only a handful of recent records. ( Extinct? Only two, possibly doubtful, records since 1970. Common Common
Twin-spot Carpet Mesotype didymata Twin-spotted Quaker Anorthoa munda Twin-spotted Wainscot Lenisa geminipuncta Uncertain Hoplodrina octogenaria Valerian Pug Eupithecia valerianata Vapourer Orgyia antiqua Varied Coronet Hadena compta Vestal Rhodometra sacraria Vine's Rustic Hoplodrina ambigua V-Hog Chloroclystis v-ata Water Carpet Lampropteryx suffumata Water Ermine Spilosoma urticae Waved Black Parascotia fuliginaria Waved Umber Menophra abruptaria Webb's Wainscot Globia sparganii White Ermine Spilosoma lubricipeda White Satin Moth Leucoma salicis White-line Dart Euxoa tritici White-marked Cerastis leucographa White-pinion Spotted Lomographa bimaculata White-point Mythimna albipuncta White-spotted Pinion Cosmia diffinis White-spotted Pug Eupithecia tripunctaria Willow Beauty Peribatodes rhomboidaria Winter Moth Operophtera brumata Wood Carpet Epirrhoe rivata Wood Tiger Parasemia plantaginis Wormwood Pug Eupithecia absinthiia	Fairly common Fairly common in suitable habitats. Uncommon. Mainly recorded from Rutland Water, but occurs elsewhere Common. Only three reliable records, from two woodland sites in east of county. Common especially in urban areas Now fairly common in gardens Formerly an uncommon migrant, now seen more frequently. Common, especially in eastern half of county Very Rare. Declined dramatically. Now only know from Clipsham Quarry Common. Fairly common Extinct. VCH & a couple of other records which need investigating Very rare. Only six records, all recent. Increasing? Common Very rare. Only a handful of recent records Common Fairly common Fairly common Fairly common Fairly rare Rare & local, restricted to the east of County Dramatic increase since mid-1980's, now common, less so in gardens Very rare migrant. Only a handful of recent records. ( Extinct? Only two, possibly doubtful, records since 1970. Common Common Common Very rare/status uncertain - all recent records need checking/verification. Extinct. Was locally common, but not recorded in last 50 years
Twin-spot Carpet Mesotype didymata Twin-spotted Quaker Anorthoa munda Twin-spotted Wainscot Lenisa geminipuncta Uncertain Hoplodrina octogenaria Valerian Pug Eupithecia valerianata Vapourer Orgyia antiqua Varied Coronet Hadena compta Vestal Rhodometra sacraria Vine's Rustic Hoplodrina ambigua V-Moth Macaria wauaria V-Pug Chloroclystis v-ata Water Carpet Lampropteryx suffumata Water Ermine Spilosoma urticae Waved Black Parascotia fuliginaria Waved Umber Menophra abruptaria Webb's Wainscot Globia sparganii White Ermine Spilosoma lubricipeda White Satin Moth Leucoma salicis White-line Dart Euxoa tritici White-marked Cerastis leucographa White-pinion Spotted Lomographa bimaculata White-spotted Pinion Cosmia diffinis White-spotted Pug Eupithecia tripunctaria Willow Beauty Peribatodes rhomboidaria Winter Moth Operophtera brumata Wood Carpet Epirrhoe rivata Wood Tiger Parasemia plantaginis Wormwood Pug Eupithecia absinthiata Yellow Horned Achlya flavicornis	Fairly common Fairly common in suitable habitats. Uncommon. Mainly recorded from Rutland Water, but occurs elsewhere Common. Only three reliable records, from two woodland sites in east of county. Common especially in urban areas Now fairly common in gardens Formerly an uncommon migrant, now seen more frequently. Common, especially in eastern half of county Very Rare. Declined dramatically. Now only know from Clipsham Quarry Common. Fairly common Extinct. VCH & a couple of other records which need investigating Very rare. Only six records, all recent. Increasing? Common Very rare. Only a handful of recent records Common Fairly common Fairly common Fairly rare Rare & local, restricted to the east of County Dramatic increase since mid-1980's, now common, less so in gardens Very rare migrant. Only a handful of recent records. ( Extinct? Only two, possibly doubtful, records since 1970. Common Common Common Common Common Common Lexinct. Was locally common, but not recorded in last 50 years Rare. Most records are of larvae found on Wormwood. Common, particularly in urban areas. Locally common
Twin-spot Carpet Mesotype didymata Twin-spotted Quaker Anorthoa munda Twin-spotted Wainscot Lenisa geminipuncta Uncertain Hoplodrina octogenaria Valerian Pug Eupithecia valerianata Vapourer Orgyia antiqua Varied Coronet Hadena compta Vestal Rhodometra sacraria Vine's Rustic Hoplodrina ambigua V-Moth Macaria wauaria V-Pug Chloroclystis v-ata Water Carpet Lampropteryx suffumata Water Ermine Spilosoma urticae Waved Black Parascotia fuliginaria Waved Umber Menophra abruptaria Webb's Wainscot Globia sparganii White Ermine Spilosoma lubricipeda White Satin Moth Leucoma salicis White-line Dart Euxoa tritici White-marked Cerastis leucographa White-pinion Spotted Lomographa bimaculata White-spotted Pinion Cosmia diffinis White-spotted Pug Eupithecia tripunctaria Willow Beauty Peribatodes rhomboidaria Winter Moth Operophtera brumata Wood Carpet Epirrhoe rivata Wood Tiger Parasemia plantaginis Wormwood Pug Eupithecia absinthiata Yellow Horned Achlya flavicornis Yellow Shell Camptogramma bilineata	Fairly common Fairly common in suitable habitats. Uncommon. Mainly recorded from Rutland Water, but occurs elsewhere Common. Only three reliable records, from two woodland sites in east of county. Common especially in urban areas Now fairly common in gardens Formerly an uncommon migrant, now seen more frequently. Common, especially in eastern half of county Very Rare. Declined dramatically. Now only know from Clipsham Quarry Common. Fairly common Extinct. VCH & a couple of other records which need investigating Very rare. Only six records, all recent. Increasing? Common Very rare. Only a handful of recent records Common Fairly common Fairly common Fairly common Fairly rare Rare & local, restricted to the east of County Dramatic increase since mid-1980's, now common, less so in gardens Very rare migrant. Only a handful of recent records. ( Extinct? Only two, possibly doubtful, records since 1970. Common Common Common Common Common Very rare/status uncertain - all recent records need checking/verification. Extinct. Was locally common, but not recorded in last 50 years Rare. Most records are of larvae found on Wormwood. Common, particularly in urban areas. Locally common
Twin-spot Carpet Mesotype didymata Twin-spotted Quaker Anorthoa munda Twin-spotted Wainscot Lenisa geminipuncta Uncertain Hoplodrina octogenaria Valerian Pug Eupithecia valerianata Vapourer Orgyia antiqua Varied Coronet Hadena compta Vestal Rhodometra sacraria Vine's Rustic Hoplodrina ambigua V-Hug Chloroclystis v-ata Water Carpet Lampropteryx suffumata Water Ermine Spilosoma urticae Waved Black Parascotia fuliginaria Waved Umber Menophra abruptaria Webb's Wainscot Globia sparganii White Ermine Spilosoma lubricipeda White Satin Moth Leucoma salicis White-line Dart Euxoa tritici White-marked Cerastis leucographa White-pinion Spotted Lomographa bimaculata White-point Mythimna albipuncta White-spotted Pinion Cosmia diffinis White-spotted Pug Eupithecia tripunctaria Willow Beauty Peribatodes rhomboidaria Winter Moth Operophtera brumata Wood Carpet Epirrhoe rivata Wood Tiger Parasemia plantaginis Wormwood Pug Eupithecia absinthiata Yellow Horned Achlya flavicornis Yellow Shell Camptogramma bilineata Yellow-barred Brindle Acasis viretata	Fairly common Fairly common in suitable habitats. Uncommon. Mainly recorded from Rutland Water, but occurs elsewhere Common. Only three reliable records, from two woodland sites in east of county. Common especially in urban areas Now fairly common in gardens Formerly an uncommon migrant, now seen more frequently. Common, especially in eastern half of county Very Rare. Declined dramatically. Now only know from Clipsham Quarry Common. Fairly common Extinct. VCH & a couple of other records which need investigating Very rare. Only six records, all recent. Increasing? Common Very rare. Only a handful of recent records Common Fairly common Fairly rare Rare & local, restricted to the east of County Dramatic increase since mid-1980's, now common, less so in gardens Very rare migrant. Only a handful of recent records. ( Extinct? Only two, possibly doubtful, records since 1970. Common Common Common Common Very rare/status uncertain - all recent records need checking/verification. Extinct. Was locally common, but not recorded in last 50 years Rare. Most records are of larvae found on Wornwood. Common Common Common Fairly common Fairly common Fairly common Fairly common
Twin-spot Carpet Mesotype didymata Twin-spotted Quaker Anorthoa munda Twin-spotted Wainscot Lenisa geminipuncta Uncertain Hoplodrina octogenaria Valerian Pug Eupithecia valerianata Vapourer Orgyia antiqua Varied Coronet Hadena compta Vestal Rhodometra sacraria Vine's Rustic Hoplodrina ambigua V-Moth Macaria wauaria V-Pug Chloroclystis v-ata Water Carpet Lampropteryx suffumata Water Ermine Spilosoma urticae Waved Black Parascotia fuliginaria Webb's Wainscot Globia sparganii White Ermine Spilosoma lubricipeda White Satin Moth Leucoma salicis White-line Dart Euxoa tritici White-marked Cerastis leucographa White-pinion Spotted Lomographa bimaculata White-point Mythimna albipuncta White-spotted Pinion Cosmia diffinis White-spotted Pug Eupithecia tripunctaria Willow Beauty Peribatodes rhomboidaria Winter Moth Operophtera brumata Wood Carpet Epirrhoe rivata Wood Tiger Parasemia plantaginis Wormwood Cucullia absinthii Wormwood Pug Eupithecia absinthiata Yellow-barred Brindle Acasis viretata Yellow-barred Brindle Acasis viretata Yellow-barred Brindle Acasis viretata	Fairly common Fairly common in suitable habitats. Uncommon. Mainly recorded from Rutland Water, but occurs elsewhere Common. Only three reliable records, from two woodland sites in east of county. Common especially in urban areas Now fairly common in gardens Formerly an uncommon migrant, now seen more frequently. Common, especially in eastern half of county Very Rare. Declined dramatically. Now only know from Clipsham Quarry Common. Fairly common Extinct. VCH & a couple of other records which need investigating Very rare. Only six records, all recent. Increasing? Common Very rare. Only a handful of recent records Common Fairly common Fairly common Fairly rare Rare & local, restricted to the east of County Dramatic increase since mid-1980's, now common, less so in gardens Very rare migrant. Only a handful of recent records. ( Extinct? Only two, possibly doubtful, records since 1970. Common Pairly common, but not recorded in last 50 years Rare. Most records are of larvae found on Wormwood. Common, particularly in urban areas. Locally common. Uncommon/under-recorded (can be common at pheromone lures)
Twin-spot Carpet Mesotype didymata Twin-spotted Quaker Anorthoa munda Twin-spotted Wainscot Lenisa geminipuncta Uncertain Hoplodrina octogenaria Valerian Pug Eupithecia valerianata Vapourer Orgyia antiqua Varied Coronet Hadena compta Vestal Rhodometra sacraria Vine's Rustic Hoplodrina ambigua V-Moth Macaria wauaria V-Pug Chloroclystis v-ata Water Carpet Lampropteryx suffumata Water Ermine Spilosoma urticae Waved Black Parascotia fuliginaria Webb's Wainscot Globia sparganii White Ermine Spilosoma lubricipeda White Satin Moth Leucoma salicis White-line Dart Euxoa tritici White-marked Cerastis leucographa White-pinion Spotted Lomographa bimaculata White-point Mythimna albipuncta White-spotted Pinion Cosmia diffinis White-spotted Pug Eupithecia tripunctaria Willow Beauty Peribatodes rhomboidaria Winter Moth Operophtera brumata Wood Carpet Epirrhoe rivata Wood Tiger Parasemia plantaginis Wormwood Pug Eupithecia absinthiata Yellow Horned Achlya flavicornis Yellow Shell Camptogramma bilineata Yellow-barred Brindle Acasis viretata	Fairly common Fairly common in suitable habitats. Uncommon. Mainly recorded from Rutland Water, but occurs elsewhere Common. Only three reliable records, from two woodland sites in east of county. Common especially in urban areas Now fairly common in gardens Formerly an uncommon migrant, now seen more frequently. Common, especially in eastern half of county Very Rare. Declined dramatically. Now only know from Clipsham Quarry Common. Fairly common Extinct. VCH & a couple of other records which need investigating Very rare. Only six records, all recent. Increasing? Common Very rare. Only a handful of recent records Common Fairly common Fairly rare Rare & local, restricted to the east of County Dramatic increase since mid-1980's, now common, less so in gardens Very rare migrant. Only a handful of recent records. ( Extinct? Only two, possibly doubtful, records since 1970. Common Common Common Common Very rare/status uncertain - all recent records need checking/verification. Extinct. Was locally common, but not recorded in last 50 years Rare. Most records are of larvae found on Wornwood. Common Common Common Fairly common Fairly common Fairly common Fairly common